

On the values of a class of analytic functions at algebraic points

Boris ADAMCZEWSKI, Yann BUGEAUD and Florian LUCA

Abstract. *We introduce a class of analytic functions of number theoretic interest, namely stammering functions. It has been recently proved that these functions take transcendental values at some algebraic points. In the present paper, we establish a general transcendence criterion that extends these results. Another aim is to underline the main difficulties arising from the use of the Schmidt Subspace Theorem in this context.*

1. Introduction

In 1844, Liouville established that the real number $\sum_{k \geq 1} 10^{-k!}$ is transcendental. With a similar method, it is nowadays an easy exercise to extend his result as follows (see [8], page 2).

Theorem L. *Let α be an algebraic number with $0 < |\alpha| < 1$. Then, the complex number $\sum_{k \geq 1} \alpha^{k!}$ is transcendental.*

In other words, the analytic function $f : z \mapsto \sum_{k \geq 1} z^{k!}$ takes transcendental values at every nonzero algebraic point in its open disc of convergence.

A similar result was obtained by Mahler [6] for a much wider class of functions f , including some classical series such as $\mathcal{F}(z) = \sum_{k \geq 0} z^{2^k}$ or the Thue–Morse function $\mathcal{T}(z) = \sum_{k \geq 0} t_k z^k$, where $t_k = 1$ (resp. $t_k = -1$), if the sum of digits of the binary expansion of k is even (resp. odd). His method was subsequently refined and generalized by several authors including Kubota, Loxton, van der Poorten, Masser and Nishioka, and the reader is referred to Nishioka’s Lecture Notes [8] for references. Since it always relies on certain functional equations satisfied by the relevant series (for instance, we have $\mathcal{F}(z^2) = \mathcal{F}(z) - z$ and $\mathcal{T}(z) = (1 - z)\mathcal{T}(z^2)$), Mahler’s method does not allow much flexibility.

The heart of the proof of Theorem L and of Mahler’s results is a lower estimate for the distance between two distinct algebraic numbers. Thanks to works by Thue, Siegel, Roth and Schmidt, the seminal result of Liouville has been considerably improved. Thus, it is not surprising that the use of the Schmidt Subspace Theorem in the present context yields a considerable improvement of Theorem L. This was recently worked out for lacunary functions by Corvaja and Zannier [5], who developed a new approach. We quote below a

2000 *Mathematics Subject Classification* : 11J81, 11J61.

particular case of Corollary 5 from [5], which is proved by means of a nontrivial application of the Schmidt Subspace Theorem.

Theorem CZ. *Let $(m_k)_{k \geq 1}$ be an increasing sequence of positive integers such that $\liminf_k m_{k+1}/m_k > 1$. Then, the function $f : z \mapsto \sum_{k \geq 1} z^{m_k}$ takes transcendental values at every algebraic point in the open unit disc.*

It is likely that Theorem CZ cannot be proved using Mahler’s method or its refinements. Furthermore, the method of [5] offers much more flexibility than Mahler’s. Unfortunately, it does not seem to apply easily to nonlacunary functions such as the Thue–Morse function \mathcal{T} .

In two earlier works [3, 1], we established new transcendence criteria for analytic functions with special combinatorial properties evaluated at the inverse of positive rational integers, Pisot and Salem numbers. These results apply to a broad class of functions including \mathcal{T} and the lacunary series occurring in Theorem CZ (see Section 2 for more details). As in [5], their proofs rest on the Schmidt Subspace Theorem. It is the aim of the present paper to investigate how our method can be applied to every algebraic point in the open disc of convergence of these analytic functions. Our main result is a general transcendence criterion that extends those of [3] and [1].

2. Stammering functions

In this section, we introduce the notion of stammering function.

Let \mathcal{A} be a countable set. The length of a word W on the alphabet \mathcal{A} , that is, the number of letters composing W , is denoted by $|W|$. For every positive integer ℓ , we write W^ℓ for the word $W \dots W$ (the concatenation of the word W repeated ℓ times). More generally, for every positive real number x , we denote by W^x the word $W^{\lceil x \rceil} W'$, where W' is the prefix of W of length $\lceil (x - \lfloor x \rfloor) |W| \rceil$. Here, and in all what follows, $\lfloor y \rfloor$ and $\lceil y \rceil$ denote, respectively, the floor and the ceiling of the real number y . Let $\mathbf{a} = (a_k)_{k \geq 0}$ be a sequence of elements from \mathcal{A} . We say that \mathbf{a} is a *stammering sequence* if \mathbf{a} is not eventually periodic and if there exist real numbers $w' \geq 0$ and $w > 1$, and two sequences of finite words $(U_n)_{n \geq 1}$, $(V_n)_{n \geq 1}$ such that:

- (i) For any $n \geq 1$, the word $U_n V_n^w$ is a prefix of the word \mathbf{a} ;
- (ii) The sequence $(|U_n|/|V_n|)_{n \geq 1}$ is bounded from above by w' ;
- (iii) The sequence $(|V_n|)_{n \geq 1}$ is strictly increasing.

Several classical sequences studied in symbolic dynamics, number theory and combinatorics on words turn out to be stammering, as pointed out in [1]. We quote below some famous examples of stammering sequences:

- Sequences generated by finite automata (these include the Thue–Morse, the Rudin–Shapiro, the Baum–Sweet, and the regular paperfolding sequences);
- Most of the morphic sequences (a morphic sequence is a letter-to-letter projection of a fixed point of an endomorphism from a finitely generated free monoid; for instance,

the Fibonacci word is the fixed point of the binary morphism σ defined by $\sigma(0) = 01$ and $\sigma(1) = 0$);

- Sturmian (or Beatty) sequences (i.e., sequences $(a_k)_{k \geq 0}$ such that there exist an irrational real number θ and a real number ρ both in $[0, 1]$ such that either $a_k = \lfloor (k+1)\theta + \rho \rfloor - \lfloor k\theta + \rho \rfloor$ for every nonnegative integer k , or $a_k = \lceil (k+1)\theta + \rho \rceil - \lceil k\theta + \rho \rceil$ for every nonnegative integer k);
- Sequences with sublinear block-complexity (these include natural codings of interval exchange transformations, billiard sequences, Arnoux–Rauzy sequences, ...);
- Characteristic sequences of lacunary sets of integers (i.e., sequences $(a_k)_{k \geq 0}$ such that $a_k = 1$ if k belongs to $\mathbf{m} = (m_\ell)_{\ell \geq 1}$ and $a_k = 0$ otherwise, where \mathbf{m} is an increasing sequence such that $\limsup_{\ell \rightarrow \infty} m_{\ell+1}/m_\ell > 1$).

We refer the reader to [1] and [4] for more details on these sequences.

Throughout the present paper, we denote by h (resp. by H) the logarithmic (resp. multiplicative) absolute height. Their definitions and main properties are given in Section 4. Furthermore, we use the symbols o , O and \ll with their usual meanings.

We say that a sequence of algebraic numbers $(a_k)_{k \geq 0}$ satisfies the growth condition (G) if either

$$h(a_0, \dots, a_n) = o(n), \quad \text{as } n \rightarrow \infty, \quad (2.1)$$

or

$$h(a_k) \ll 1. \quad (2.2)$$

Obviously, by (2.2), every bounded sequence of rational integers satisfies the growth condition (G).

With a stammering sequence $\mathbf{a} = (a_k)_{k \geq 0}$ of algebraic numbers in a number field \mathbf{L} that satisfies the growth condition (G), we associate the analytic function $f_{\mathbf{a}}$ defined on the open unit disc by

$$f_{\mathbf{a}}(z) = \sum_{k=0}^{+\infty} a_k z^k. \quad (2.3)$$

Such function is called a *stammering function*. This notation will be kept throughout the entire paper. We note that all the series mentioned in Section 1 are stammering ones.

3. Results

We are interested in values of stammering functions at algebraic points. In view of the results mentioned previously, it is likely that such functions essentially take transcendental values at algebraic points. More precisely, we can expect the following picture: given a stammering function $f_{\mathbf{a}}$ as in (2.3), then $f_{\mathbf{a}}(\alpha)$ is transcendental for all but finitely many algebraic numbers α lying in the open disc of convergence of $f_{\mathbf{a}}$, and the algebraic values of $f_{\mathbf{a}}$ belong to the number field $\mathbf{L}(\alpha)$.

A first result towards this problem was proved in [3].

Theorem ABL. *Let $b \geq 2$ be a rational integer. If $\mathbf{a} = (a_k)_{k \geq 0}$ is a stammering sequence on the set $\{0, 1, \dots, b-1\}$, then the stammering function $f_{\mathbf{a}}$ takes transcendental values at the inverse of every positive rational integer.*

As worked out in [1], this transcendence criterion is powerful enough to confirm the Cobham–Loxton–van der Poorten conjecture claiming that the b -adic expansion of an algebraic irrational real number cannot be generated by a finite automaton. The above is a result that seems to escape to Mahler’s method.

Subsequently, Theorem ABL was extended in [1] in the following way. Recall that a Pisot (resp. Salem) number is a real algebraic integer > 1 , whose complex conjugates lie inside the open unit disc (resp. inside the closed unit disc, with at least one of them on the unit circle).

Theorem AB. *Let $\mathbf{a} = (a_k)_{k \geq 0}$ be a stammering, bounded sequence of rational integers. If β is a Pisot or a Salem number, then $f_{\mathbf{a}}(1/\beta)$ is either transcendental, or belongs to the number field $\mathbf{Q}(\beta)$.*

Note that we cannot avoid the possibility that $f_{\mathbf{a}}(1/\beta)$ belongs to $\mathbf{Q}(\beta)$. Indeed, let $\beta = (\sqrt{5}+1)/2$ be the Golden Ratio. Then, starting with a periodic sequence $\mathbf{a} = (a_k)_{k \geq 1}$, we can use the fact that $1 = 1/\beta + 1/\beta^2$ to construct (using only local perturbation of the sequence \mathbf{a}) a stammering sequence $\mathbf{a}' = (a'_k)_{k \geq 0}$ such that

$$\mathbf{Q}(\beta) \ni \sum_{k=0}^{+\infty} \frac{a_k}{\beta^k} = \sum_{k=0}^{+\infty} \frac{a'_k}{\beta^k}.$$

The proof of Theorem AB rests on a p -adic version due to Schlickewei [9] of the Schmidt Subspace Theorem. The crucial point to apply the Subspace Theorem is that the linear form $f_{\mathbf{a}}(\alpha)X - f_{\mathbf{a}}(\alpha)Y - Z$ takes small values at many algebraic points when $\alpha := 1/\beta$ is the inverse of a Pisot or a Salem number. Unfortunately, when α is an arbitrary complex algebraic number lying in the open unit disc, the values taken by this linear form are no longer small enough with respect to the height of the relevant algebraic points. Notice also that a similar difficulty explains the condition imposed on the constant L in Theorem 3 of Corvaja and Zannier [5].

If $|\cdot|_j$ is an Archimedean absolute value, given by the embedding σ_j , such that $|\alpha|_j < 1$, then we can consider the complex number $f_{\mathbf{a},j}(\alpha)$ defined by

$$f_{\mathbf{a},j}(\alpha) := \lim_{N \rightarrow +\infty} \sum_{k=0}^N a_k \sigma_j(\alpha^k).$$

Again, if \mathfrak{p} is a finite place such that $|\alpha|_{\mathfrak{p}} < 1$, then we can consider in the \mathfrak{p} -adic completion of the field $\mathbf{Q}(\beta)$ the number $f_{\mathbf{a},\mathfrak{p}}(\alpha)$ defined by

$$f_{\mathbf{a},\mathfrak{p}}(\alpha) := \lim_{N \rightarrow +\infty} \sum_{k=0}^N a_k \alpha^k.$$

The main novelty of the present paper is to remark that all the linear forms $f_{\mathbf{a},j}(\alpha)X - f_{\mathbf{a},j}(\alpha)Y - Z$ and $f_{\mathbf{a},\mathfrak{p}}(\alpha)X - f_{\mathbf{a},\mathfrak{p}}(\alpha)Y - Z$ also take small values at many algebraic points constructed in the same way as in the proof of Theorem AB. Actually, when evaluated at such points the product of all linear forms we consider is just small enough to apply the Subspace Theorem.

Our main result is the following extension of Theorem AB to any nonzero complex algebraic number lying in the open unit disc.

Theorem 1. *Let $\mathbf{a} = (a_k)_{k \geq 0}$ be a stammering sequence of algebraic numbers in a number field \mathbf{L} and satisfying the growth condition (G). Let α be a nonzero algebraic number lying in the open unit disc and assume that $\alpha_1 = \alpha, \alpha_2, \dots, \alpha_\ell$ are all its complex conjugates of modulus < 1 . Let $\mathfrak{p}_1, \dots, \mathfrak{p}_{\ell'}$ be all the prime ideals in $\mathbf{L}(\alpha)$ such that $|\alpha|_{\mathfrak{p}_j} < 1$. Then, either (at least) one among the numbers $f_{\mathbf{a},1}(\alpha) := f_{\mathbf{a}}(\alpha), f_{\mathbf{a},2}(\alpha), \dots, f_{\mathbf{a},\ell}(\alpha), f_{\mathbf{a},\mathfrak{p}_1}(\alpha), \dots, f_{\mathbf{a},\mathfrak{p}_{\ell'}}(\alpha)$ is transcendental, or $f_{\mathbf{a},j}(\alpha)$ belongs to $\mathbf{L}(\alpha_j)$ for $j = 1, \dots, \ell$. Furthermore, in this last case letting σ_j be some automorphism of \mathbf{K} sending α to α_j , then the relation $f_{\mathbf{a},j}(\alpha) = \sigma_j(f_{\mathbf{a}}(\alpha))$ holds for all $j = 1, \dots, \ell$.*

We stress that Theorem 1 plainly includes Theorems ABL and AB. The proof of Theorem 1 also rests on the p -adic version of the Schmidt Subspace Theorem given in [9]. The main interest of Theorem 1 is that it applies to every algebraic value of every stammering function. However, the price to pay for this (i.e., the introduction of new linear forms) produces a weaker conclusion than the one expected. We mention that another way to proceed, worked out in [2], yields a conclusion as strong as in Theorem AB, but the price to pay is then a strengthening of the stammering condition on \mathbf{a} .

We give now two straightforward consequences of Theorem 1.

Corollary 1. *Let $\mathbf{a} = (a_k)_{k \geq 0}$ be a stammering sequence of algebraic numbers in a number field \mathbf{L} and satisfying the growth condition (G). Let β be an algebraic integer such that $|\beta| > 1$ and assume that $\beta_1 = \beta, \beta_2, \dots, \beta_\ell$ are all its complex conjugates of modulus > 1 . If all the complex numbers*

$$\gamma_j := \sum_{k=0}^{+\infty} \frac{a_k}{\beta_j^k}, \quad 1 \leq j \leq \ell,$$

are algebraic, then γ_j belongs to the number field $\mathbf{L}(\beta_j)$, for $j = 1, \dots, \ell$.

Corollary 2. *Let $\mathbf{a} = (a_k)_{k \geq 0}$ be a bounded, stammering sequence of rational integers. Let p and q be positive coprime integers with $q > p$ and p prime. If both the real number*

$$\gamma := \sum_{k=0}^{+\infty} a_k \left(\frac{p}{q}\right)^k$$

and the p -adic number

$$\sum_{k=0}^{+\infty} a_k \left(\frac{p}{q}\right)^k$$

are algebraic, then γ is rational.

We end this section with an application of Theorem 1 to the expansion of algebraic numbers in a complex quadratic integer base b , which in turn has an amusing application to the transcendence of alternating stammering numbers.

Theorem 2. *Let b be an algebraic integer in an imaginary quadratic field which is not a root of unity. Let $\mathbf{a} = (a_k)_{k \geq 0}$ be a stammering sequence with values in the set $\{0, 1, \dots, |b|^2 - 1\}$. Then, the number*

$$\sum_{k=0}^{+\infty} a_k/b^k$$

is transcendental.

Corollary 3. *If $b > 1$ is a positive integer and $\mathbf{a} = (a_k)_{k \geq 0}$ is a stammering sequence taking its values in the set $\{0, 1, \dots, b - 1\}$, then the number*

$$\sum_{k=0}^{+\infty} (-1)^k a_k/b^k$$

is transcendental.

4. The auxiliary result

Our main auxiliary tool is a version of the \mathfrak{P} -adic Thue–Siegel–Roth–Schmidt theorem proved by Schlickewei [9]. We make use of the following notation (see also [7], pp. 182–183).

Let p be a prime number. Let \mathbf{Q}_p be the completion of \mathbf{Q} at p and \mathbf{C}_p the completion of the algebraic closure of \mathbf{Q}_p for the absolute value $|\cdot|_p$.

Let \mathbf{K} be a number field of degree d . Write the ideal (p) in $\mathcal{O}_{\mathbf{K}}$ as

$$(p) = \prod_{j=1}^{\pi(p)} \mathfrak{p}_j^{e(\mathfrak{p}_j)},$$

where $\mathfrak{p}_1, \dots, \mathfrak{p}_{\pi(p)}$ are distinct prime ideals in $\mathcal{O}_{\mathbf{K}}$. For a nonzero element α in \mathbf{K} , set

$$|\alpha|_{\mathfrak{p}_j} = p^{-h_j/e(\mathfrak{p}_j)},$$

where h_j is the largest integer such that \mathfrak{p}_j is coprime with the fractional ideal $\mathfrak{p}_j^{h_j} \times (\alpha)$. For $j = 1, \dots, \pi(p)$, let $d(\mathfrak{p}_j)$ denote the degree of the completion $\mathbf{K}_{\mathfrak{p}_j}$ of \mathbf{K} with respect to \mathfrak{p}_j . Let r be the unit rank of \mathbf{K} . For $j = 1, \dots, r + 1$, let d_j equal to 1 (resp. 2) if the embedding of \mathbf{K} associated with the absolute value $|\cdot|_j$ is real (resp. complex).

For any nonzero α in \mathbf{K} , we have the product formula

$$\prod_{j=1}^{r+1} |\alpha|_j^{d_j} \cdot \prod_{\mathfrak{p}} |\alpha|_{\mathfrak{p}}^{d(\mathfrak{p})} = 1.$$

The field height $H_{\mathbf{K}}$ of $\underline{\beta} = (\beta_0, \dots, \beta_m) \in \mathbf{K}^{m+1}$ is defined by

$$H_{\mathbf{K}}(\underline{\beta}) = \prod_{j=1}^{r+1} \max\{|\beta_0|_j^{d_j}, \dots, |\beta_m|_j^{d_j}\} \prod_{\mathfrak{p}} \max\{|\beta_0|_{\mathfrak{p}}^{d(\mathfrak{p})}, \dots, |\beta_m|_{\mathfrak{p}}^{d(\mathfrak{p})}\},$$

where the second product is taken over all inequivalent nonArchimedean primes of \mathbf{K} .

Let $p_0 = \infty$ denote the Archimedean prime in \mathbf{Q} . Denote by $\mathbf{K}_1^{(0)}, \dots, \mathbf{K}_{r+1}^{(0)}$ the r pairwise nonequivalent isomorphic embeddings of \mathbf{K} into \mathbf{C} . For simplicity, set $\pi(p_0) = r + 1$. Let $\{p_1, \dots, p_t\}$ be a finite set of rational prime numbers. For $i = 1, \dots, t$, denote by $\mathbf{K}_1^{(i)}, \dots, \mathbf{K}_{\pi(p_i)}^{(i)}$ the $\pi(p_i)$ pairwise nonequivalent isomorphic embeddings of \mathbf{K} into \mathbf{C}_{p_i} . Furthermore, we denote the images of an $(m + 1)$ -tuple $\underline{\beta} = (\beta_0, \dots, \beta_m) \in \mathbf{K}^{m+1}$ under these embeddings as follows: for $i = 0, \dots, t$ and for $j = 1, \dots, \pi(p_i)$, we write $\underline{\beta}_j^{(i)} = (\beta_{0j}^{(i)}, \dots, \beta_{mj}^{(i)}) \in \mathbf{K}_j^{(i)m+1}$.

Furthermore, for $j = 1, \dots, r + 1$, set

$$|\underline{\beta}_j^{(0)}| = \max\{|\beta_{0j}^{(0)}|, \dots, |\beta_{mj}^{(0)}|\},$$

and, for $i = 1, \dots, t$ and $j = 1, \dots, \pi(p_i)$, set

$$|\underline{\beta}_j^{(i)}|_{\mathfrak{p}_{ij}} = \max\{|\beta_{0j}^{(i)}|_{\mathfrak{p}_{ij}}, \dots, |\beta_{mj}^{(i)}|_{\mathfrak{p}_{ij}}\}.$$

With these notations, Theorem 2.1 of [9] combined with Lemma 1F, page 178, of [10] yields the following version of the Subspace Theorem, that is the key tool for the proof of our Theorem 1.

Theorem S. *For every $i = 0, \dots, t$ and every $j = 1, \dots, \pi(p_i)$, let $L_{0j}^{(i)}, \dots, L_{mj}^{(i)}$ be $m + 1$ linearly independent linear forms in $m + 1$ variables with algebraic coefficients from the field \mathbf{C}_{p_i} . Then, for any $\varepsilon > 0$, all the solutions $\underline{\beta} \in \mathbf{P}^m(\mathbf{K})$ to the inequality*

$$\prod_{j=1}^{r+1} \prod_{k=0}^m \frac{|L_{kj}^{(0)}(\underline{\beta}_j^{(0)})|^{d_j}}{|\underline{\beta}_j^{(0)}|^{d_j}} \cdot \prod_{i=1}^t \prod_{j=1}^{\pi(p_i)} \prod_{k=0}^m \frac{|L_{kj}^{(i)}(\underline{\beta}_j^{(i)})|_{\mathfrak{p}_{ij}}^{d(\mathfrak{p}_{ij})}}{|\underline{\beta}_j^{(i)}|_{\mathfrak{p}_{ij}}^{d(\mathfrak{p}_{ij})}} \leq H_{\mathbf{K}}(\underline{\beta})^{-m-1-\varepsilon} \quad (4.1)$$

lie in finitely many proper subspaces.

We end this section with a few words on heights. Keep the above notation and let β be in \mathbf{K} . The (multiplicative) absolute height of β is defined by

$$H(\beta) := (H_{\mathbf{K}}((1, \beta)))^{1/[\mathbf{K}:\mathbf{Q}]},$$

and we set

$$h(\beta) = \log H(\beta).$$

We refer to Chapter 3 of [11] for classical results on these heights.

We just mention that, for any algebraic numbers a_0, \dots, a_n, β in \mathbf{K} and for any place ν in \mathbf{K} we have

$$\begin{aligned} |a_0 + a_1\beta + \dots + a_n\beta^n|_\nu &\leq \max\{1, |n+1|_\nu\} \times \max\{|a_0|_\nu, |a_1\beta|_\nu, \dots, |a_n\beta^n|_\nu\} \\ &\leq \max\{1, |n+1|_\nu\} \times \max\{|a_0|_\nu, \dots, |a_n|_\nu\} \\ &\quad \times \max\{1, |\beta|_\nu\}^n, \end{aligned} \tag{4.2}$$

and

$$h(a_0, \dots, a_n) := \frac{\log(H_{\mathbf{K}}((a_0, \dots, a_n)))}{[\mathbf{K} : \mathbf{Q}]}.$$
 \tag{4.3}

5. Proof of Theorem 1

This section is devoted to the proof of Theorem 1.

Proof of Theorem 1. Consider a stammering sequence $\mathbf{a} = (a_k)_{k \geq 0}$ of algebraic numbers in a number field \mathbf{L} that satisfies the growth condition (G). We assume that the parameters w and w' are fixed, as well as the sequences $(U_n)_{n \geq 1}$ and $(V_n)_{n \geq 1}$ occurring in the definition of a stammering sequence. Set also $r_n = |U_n|$ and $s_n = |V_n|$, for every $n \geq 1$. Observe that s_n tends to infinity monotonically with n .

We fix a complex algebraic number α in the open unit disc and set $\beta := 1/\alpha$. We point out that $|\beta| > 1$. Set $\mathbf{K} := \mathbf{L}(\alpha) = \mathbf{L}(\beta)$ and $\delta = [\mathbf{K} : \mathbf{Q}]/[\mathbf{Q}(\beta) : \mathbf{Q}]$. Assume that the conjugates of β are numbered in such a way that $\beta = \beta_1, \dots, \beta_\ell$ is a subset of the complex conjugates of β of modulus > 1 containing exactly one of the two elements of each pair of conjugates.

For any $j = 1, \dots, \ell$, set

$$f_{\mathbf{a},j}(\alpha) := f_{\mathbf{a}}(\alpha_j) = \sum_{k=0}^{+\infty} a_k \alpha_j^k = \sum_{k=0}^{+\infty} \frac{a_k}{\beta_j^k}.$$

Let $\mathfrak{p}_1, \dots, \mathfrak{p}_{\ell'}$ be the prime ideals \mathfrak{p} in \mathbf{K} with $|\alpha|_{\mathfrak{p}} < 1$ (i.e., with $|\beta|_{\mathfrak{p}} > 1$). For $j = 1, \dots, \ell'$, let $f_{\mathbf{a},\mathfrak{p}_j}(\alpha)$ in $\mathbf{K}_{\mathfrak{p}_j}$ be defined by

$$f_{\mathbf{a},\mathfrak{p}_j}(\alpha) := \lim_{N \rightarrow +\infty} \sum_{k=0}^N \frac{a_k}{\beta^k}.$$

The key fact is the observation that the $f_{\mathbf{a},j}(\alpha)$'s and the $f_{\mathbf{a},\mathfrak{p}_j}(\alpha)$'s defined above admit infinitely many good approximants in the fields \mathbf{K} and $\mathbf{K}_{\mathfrak{p}_j}$, respectively. These

approximants are obtained by truncating their associated series and completing them by periodicity. Precisely, for every positive integer n , we define the sequence $(b_k^{(n)})_{k \geq 0}$ by

$$\begin{aligned} b_h^{(n)} &= a_h \quad \text{for } 0 \leq h \leq r_n + s_n - 1, \\ b_{r_n+h+j s_n}^{(n)} &= a_{r_n+h} \quad \text{for } 0 \leq h \leq s_n - 1 \text{ and } j \geq 0. \end{aligned}$$

The sequence $(b_k^{(n)})_{k \geq 0}$ is eventually periodic, with preperiod U_n and with period V_n . For $j = 1, \dots, \ell$, set

$$\alpha_j^{(n)} = \sum_{k=0}^{+\infty} \frac{b_k^{(n)}}{\beta_j^k}.$$

Since $(a_k)_{k \geq 0}$ satisfies the growth condition (G) , we have $|a_k - b_k^{(n)}| \ll c^k$ for every real number $c > 1$. Throughout the proof of the theorem, all the constants implied by \ll may depend on c , but they are independent of n . We thus observe that

$$|f_{\mathbf{a},j}(\alpha) - \alpha_j^{(n)}| = \left| \sum_{k=r_n+\lceil ws_n \rceil}^{+\infty} \frac{a_k - b_k^{(n)}}{\beta_j^k} \right| \ll \frac{c^{r_n+\lceil ws_n \rceil}}{|\beta_j|^{r_n+\lceil ws_n \rceil}}, \quad (5.1)$$

for every real number $c > 1$. Likewise, for $j = 1, \dots, \ell'$, define $\alpha_{\mathbf{p}_j}^{(n)}$ in $\mathbf{K}_{\mathbf{p}_j}$ by

$$\alpha_{\mathbf{p}_j}^{(n)} = \lim_{N \rightarrow +\infty} \sum_{k=0}^N \frac{b_k^{(n)}}{\beta^k},$$

and observe that

$$|f_{\mathbf{a},\mathbf{p}_j}(\alpha) - \alpha_{\mathbf{p}_j}^{(n)}|_{\mathbf{p}_j} \ll \frac{c^{r_n+\lceil ws_n \rceil}}{|\beta|_{\mathbf{p}_j}^{r_n+\lceil ws_n \rceil}}, \quad (5.2)$$

for every real number $c > 1$.

Lemma 1. *For every integer n , set*

$$P_n(X) = \sum_{k=0}^{r_n-1} a_k X^{r_n-k} (X^{s_n} - 1) + \sum_{k=0}^{s_n-1} a_{r_n+k} X^{s_n-k}.$$

Then, for $j = 1, \dots, \ell$, we have

$$\alpha_j^{(n)} = \frac{P_n(\beta_j)}{\beta_j^{r_n} (\beta_j^{s_n} - 1)}.$$

Proof. This is an easy verification, as in Lemma 1 from [1]. □

Now, we define the systems of linear forms to which we will apply Theorem S. Let us assume that $f_{\mathbf{a},1}(\alpha) := f_{\mathbf{a}}(\alpha), \dots, f_{\mathbf{a},\ell}(\alpha)$ and $f_{\mathbf{a},\mathbf{p}_1}(\alpha), \dots, f_{\mathbf{a},\mathbf{p}_{\ell'}}(\alpha)$ are all algebraic.

For $j = 1, \dots, \ell$, consider the linear forms

$$L_{0j}^{(0)} = X, \quad L_{1j}^{(0)} = Y, \quad L_{2j}^{(0)} = f_{\mathbf{a},j}(\alpha)X - f_{\mathbf{a},j}(\alpha)Y - Z.$$

For the remaining Archimedean absolute values, that is, for $j = \ell + 1, \dots, r + 1$, where r denotes the unit rank of $\mathbf{Q}(\beta)$, take simply

$$L_{0j}^{(0)} = X, \quad L_{1j}^{(0)} = Y, \quad L_{2j}^{(0)} = Z.$$

Let p_1, \dots, p_t be the prime numbers p such that there exists an ideal \mathfrak{p} above p with $|\beta|_{\mathfrak{p}} \neq 1$. Each of the prime ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_{\ell'}$ divides one of the p_i 's. For $j = 1, \dots, \ell'$, let p_i be the rational prime above \mathfrak{p}_j and consider the linear forms

$$L_{0j}^{(i)} = X, \quad L_{1j}^{(i)} = Y, \quad L_{2j}^{(i)} = f_{\mathbf{a},\mathfrak{p}_j}(\alpha)X - f_{\mathbf{a},\mathfrak{p}_j}(\alpha)Y - Z.$$

For all the other prime ideals \mathfrak{p} below one of the p_i 's, we consider the linear forms

$$L_{0j}^{(i)} = X, \quad L_{1j}^{(i)} = Y, \quad L_{2j}^{(i)} = Z.$$

We evaluate the product of these linear forms at the algebraic points $\underline{\beta}_j^{(i)}$, where

$$\underline{\beta} = (\beta^{r_n+s_n}, \beta^{r_n}, P_n(\beta)).$$

Obviously, $\underline{\beta}$ and the $\underline{\beta}_j^{(i)}$'s depend on n ; however, we choose not to indicate this dependence for sake of simplicity.

We first establish an upper bound for the height of $\underline{\beta}$.

Lemma 2. *For every real number $c > 1$, we have*

$$H_{\mathbf{K}}(\underline{\beta}) \ll \left(c \prod_{j=1}^{\ell} |\beta_j|^{d_j} \cdot \prod_{j=1}^{\ell'} |\beta|_{\mathfrak{p}_j}^{d_{\mathfrak{p}_j}} \right)^{r_n+s_n},$$

where the constant implied by the Vinogradov symbol \ll is independent of n .

Proof. Let $c > 1$ be arbitrary. By definition,

$$\begin{aligned} H_{\mathbf{K}}(\underline{\beta}) &= \prod_{j=1}^{r+1} \max\{|\beta_j^{r_n+s_n}|^{d_j}, |\beta_j^{r_n}|^{d_j}, |P_n(\beta_j)|^{d_j}\} \\ &\quad \cdot \prod_{\mathfrak{p}} \max\{|\beta^{r_n+s_n}|_{\mathfrak{p}}^{d(\mathfrak{p})}, |\beta^{r_n}|_{\mathfrak{p}}^{d(\mathfrak{p})}, |P_n(\beta)|_{\mathfrak{p}}^{d(\mathfrak{p})}\}. \end{aligned}$$

On the one hand, we infer from the fact that \mathbf{a} satisfies the growth condition (G) and from Lemma 1 that

$$\begin{aligned} \prod_{j=1}^{r+1} \max\{|\beta_j^{r_n+s_n}|^{d_j}, |\beta_j^{r_n}|^{d_j}, |P_n(\beta_j)|^{d_j}\} &\ll \left(\prod_{j=1}^{\ell} |\beta_j|^{d_j} \right)^{r_n+s_n} \cdot \prod_{j=\ell+1}^{r+1} \max\{1, |P_n(\beta_j)|^{d_j}\} \\ &\ll \left(\prod_{j=1}^{\ell} |\beta_j|^{d_j} \right)^{r_n+s_n} c^{(r_n+s_n)/2}. \end{aligned}$$

On the other hand, we have

$$\prod_{\mathfrak{p}} \max\{|\beta^{r_n+s_n}|_{\mathfrak{p}}^{d(\mathfrak{p})}, |\beta^{r_n}|_{\mathfrak{p}}^{d(\mathfrak{p})}, |P_n(\beta)|_{\mathfrak{p}}^{d(\mathfrak{p})}\} \ll \left(\prod_{j=1}^{\ell'} |\beta|_{\mathfrak{p}_j}^{d(\mathfrak{p}_j)} \right)^{r_n+s_n} \cdot \prod_{\mathfrak{p} \in \mathcal{P}} |P_n(\beta)|_{\mathfrak{p}}^{d(\mathfrak{p})},$$

where the latter product is taken over the set \mathcal{P} of all the prime ideals \mathfrak{p} that do not divide $\mathfrak{p}_1 \dots \mathfrak{p}_{\ell'}$ and are such that $|P_n(\beta)|_{\mathfrak{p}} > 1$. Furthermore, we have

$$\prod_{\mathfrak{p} \in \mathcal{P}} |P_n(\beta)|_{\mathfrak{p}}^{d(\mathfrak{p})} \ll c^{(r_n+s_n)/2}. \quad (5.3)$$

Indeed, we infer from (2.1), inequality (4.2), Lemma 1 and inequality (4.3) that

$$\begin{aligned} \prod_{\mathfrak{p} \in \mathcal{P}} |P_n(\beta)|_{\mathfrak{p}}^{d(\mathfrak{p})} &\leq \prod_{\mathfrak{p} \in \mathcal{P}} (\max\{1, |r_n + s_n|_{\mathfrak{p}}^{d(\mathfrak{p})}\} \cdot \max\{|a_0|_{\mathfrak{p}}^{d(\mathfrak{p})}, \dots, |a_{r_n+s_n-1}|_{\mathfrak{p}}^{d(\mathfrak{p})}\}) \\ &\leq (r_n + s_n) H_{\mathbf{K}}(a_0, \dots, a_{r_n+s_n-1}) \\ &\ll (r_n + s_n) \exp\{[\mathbf{K} : \mathbf{Q}] \cdot h(a_0, \dots, a_{r_n+s_n-1})\} \ll c^{(r_n+s_n)/2}. \end{aligned}$$

The same upper bound holds under the assumption (2.2) since, in this case, the product

$$\prod_{\mathfrak{p} \in \mathcal{P}} \max\{|a_0|_{\mathfrak{p}}^{d(\mathfrak{p})}, \dots, |a_{r_n+s_n-1}|_{\mathfrak{p}}^{d(\mathfrak{p})}\}$$

is bounded by a constant independent of n . This ends the proof. \square

Observe that our choice of p_1, \dots, p_t implies that $|\underline{\beta}|_{\mathfrak{p}} = 1$ for any prime \mathfrak{p} which does not lie above one of the p_i 's. Consequently, we get

$$\prod_{j=1}^{r+1} |\underline{\beta}_j^{(0)}|^{d_j} \cdot \prod_{i=1}^t \prod_{j=1}^{\pi(p_i)} |\underline{\beta}_j^{(i)}|_{\mathfrak{p}_{ij}}^{d(\mathfrak{p}_{ij})} = H_{\mathbf{K}}(\underline{\beta}) \cdot \prod_{\mathfrak{p} \in \mathcal{P}'} |P_n(\beta)|_{\mathfrak{p}}^{-d(\mathfrak{p})}, \quad (5.4)$$

where the latter product is taken over the set \mathcal{P}' of all the prime ideals \mathfrak{p} that do not divide $p_1 \dots p_t$ and are such that $|P_n(\beta)|_{\mathfrak{p}} > 1$. Obviously, we infer from (5.3) that

$$\prod_{\mathfrak{p} \in \mathcal{P}'} |P_n(\beta)|_{\mathfrak{p}}^{d(\mathfrak{p})} \leq \prod_{\mathfrak{p} \in \mathcal{P}} |P_n(\beta)|_{\mathfrak{p}}^{d(\mathfrak{p})} \ll c^{(r_n+s_n)/2}, \quad (5.5)$$

for every real number $c > 1$.

By the product formula, we have

$$\prod_{j=1}^{r+1} |L_{kj}^{(0)}(\underline{\beta}_j^{(0)})|^{\delta d_j} \cdot \prod_{i=1}^t \prod_{j=1}^{\pi(p_i)} |L_{kj}^{(i)}(\underline{\beta}_j^{(i)})|_{\mathfrak{p}_{ij}}^{d(\mathfrak{p}_{ij})} = 1, \quad (5.6)$$

for $k = 0, 1$.

It remains to evaluate the expressions $|L_{2j}^{(i)}(\underline{\beta}^{(i)})|$. By (5.1), the Archimedean places yield a contribution of at most

$$\prod_{j=1}^{r+1} |L_{2j}^{(0)}(\underline{\beta}^{(0)})|^{\delta d_j} \ll c_0^{r_n+s_n} \prod_{j=1}^{\ell} |\beta_j|^{-\delta d_j(w-1)s_n}, \quad (5.7)$$

for a certain real number $c_0 > 1$ that will be later on selected to be sufficiently close to 1. By (5.2), the contribution of the nonArchimedean places yields

$$\prod_{i=1}^t \prod_{j=1}^{\pi(p_i)} |L_{2j}^{(i)}(\underline{\beta}^{(i)})|_{\mathfrak{p}_{ij}}^{d(\mathfrak{p}_{ij})} \ll c_0^{r_n+s_n} \prod_{j=1}^{\ell'} |\beta|_{\mathfrak{p}_j}^{-(w-1)s_n d(\mathfrak{p}_j)}. \quad (5.8)$$

Consequently, choosing $c = c_0^{1/3}$ in (5.5), we infer from (5.4), (5.6), (5.7) and (5.8) that the product

$$\Pi := \prod_{j=1}^{r+1} \prod_{k=0}^2 \frac{|L_{kj}^{(0)}(\underline{\beta}^{(0)})|^{d_j}}{|\underline{\beta}_j^{(0)}|^{d_j}} \cdot \prod_{i=1}^t \prod_{j=1}^{\pi(p_i)} \prod_{k=0}^2 \frac{|L_{kj}^{(i)}(\underline{\beta}^{(i)})|_{\mathfrak{p}_{ij}}^{d(\mathfrak{p}_{ij})}}{|\underline{\beta}_j^{(i)}|_{\mathfrak{p}_{ij}}^{d(\mathfrak{p}_{ij})}}$$

of all the linear forms is

$$\ll H_{\mathbf{K}}(\underline{\beta})^{-3} c_0^{3(r_n+s_n)} \left(\prod_{|\beta_j|>1} |\beta_j|^{-\delta d_j(w-1)s_n} \cdot \prod_{j=1}^{\ell'} |\beta|_{\mathfrak{p}_j}^{-(w-1)s_n d(\mathfrak{p}_j)} \right).$$

Since $w > 1$, we infer from condition (ii) in the definition of a stammering sequence that there exists a positive real number η such that

$$2\eta(r_n + s_n) < (w - 1)s_n,$$

for every large positive integer n . Now, we choose c_0 close enough to 1 to ensure that $c_1 := H_{\mathbf{K}}(\underline{\beta})/c_0^{3/\eta}$ satisfies $c_1 > 1$. Since $\delta > 1$, we obtain

$$\begin{aligned} \Pi &\ll H_{\mathbf{K}}(\underline{\beta})^{-3} c_0^{3(r_n+s_n)} \left(\prod_{j=1}^{\ell} |\beta_j|^{d_j} \cdot \prod_{j=1}^{\ell'} |\beta|_{\mathfrak{p}_j}^{d(\mathfrak{p}_j)} \right)^{-2\eta(r_n+s_n)} \\ &\ll H_{\mathbf{K}}(\underline{\beta})^{-3} (c_0^{-3/\eta} H_{\mathbf{K}}(\underline{\beta}))^{-\eta(r_n+s_n)} \left(\prod_{j=1}^{\ell} |\beta_j|^{d_j} \cdot \prod_{j=1}^{\ell'} |\beta|_{\mathfrak{p}_j}^{d(\mathfrak{p}_j)} \right)^{-\eta(r_n+s_n)} \\ &\ll H_{\mathbf{K}}(\underline{\beta})^{-3} \left(c_1 \cdot \prod_{j=1}^{\ell} |\beta_j|^{d_j} \cdot \prod_{j=1}^{\ell'} |\beta|_{\mathfrak{p}_j}^{d(\mathfrak{p}_j)} \right)^{-\eta(r_n+s_n)}. \end{aligned}$$

It thus follows from Lemma 2 that

$$\Pi \ll H_{\mathbf{K}}(\beta)^{-3-\eta}.$$

This shows that we have produced infinitely many algebraic solutions to inequality (4.1). Applying Theorem S, we reach the conclusion that there exists a nonzero triple (z_1, z_2, z_3) in \mathbf{K}^3 and an infinite set \mathcal{N} of positive integers, such that

$$z_1\beta^{r_n+s_n} + z_2\beta^{r_n} + z_3P_n(\beta) = 0, \quad (5.9)$$

for all $n \in \mathcal{N}$. Since s_n tends to infinity with n , the number $z_3 \in \mathbf{K}$ is nonzero. If $z_1 \neq 0$, then dividing (5.9) by $\beta^{r_n+s_n}$ and letting n tend to infinity along \mathcal{N} , we get that

$$\lim_{n \rightarrow +\infty} \frac{P_n(\beta)}{\beta^{r_n+s_n}} = f_{\mathbf{a}}(\alpha)$$

belongs to \mathbf{K} . If $z_1 = 0$, we divide (5.9) by β^{r_n} and argue similarly to reach the same conclusion.

Given any integer j with $2 \leq j \leq \ell$, we can replace α by α_j all along the above proof, and thus derive that $f_{\mathbf{a},j}(\alpha)$ belongs to $\mathbf{L}(\alpha_j)$. The last statement about $f_{\mathbf{a},j}(\alpha) = \sigma_j(f_{\mathbf{a}}(\alpha))$ follows immediately by conjugating relation (5.9) by σ_j for all $n \in \mathcal{N}$ and then arguing as above.

This concludes the proof of our Theorem 1. □

6. Proofs of Theorem 2 and Corollary 3

This section is devoted to the proofs of Theorem 2 and Corollary 3.

Proof of Theorem 2. We may assume that $a_0 \neq 0$, since otherwise we may eliminate a_0 and reindex the sequence (clearly, the new sequence obtained in this way is still stammering).

Set

$$\alpha = \sum_{k=0}^{\infty} \frac{a_k}{b^k}.$$

Assume that $\alpha \in \overline{\mathbf{Q}}$. Since the complex conjugation is continuous, we get that

$$\bar{\alpha} = \sum_{k=0}^{\infty} \frac{a_k}{\bar{b}^k}.$$

Hence, $\bar{\alpha} \in \overline{\mathbf{Q}}$. By Theorem 1, we have that $\alpha \in \mathbf{Q}[b]$. In fact, Theorem 1 shows that more is true, namely (we keep the notation used in the proof of this theorem) that there exist

a nonzero triple (z_1, z_2, z_3) in $\mathbf{K}^3 = \mathbf{Q}[b]^3$ and an infinite set of positive integers \mathcal{N}_1 such that

$$z_1 b^{r_n+s_n} - z_2 b^{r_n} - z_3 P_n(b) = 0 \quad (6.1)$$

holds for every $n \in \mathcal{N}_1$. Here,

$$P_n(X) = \sum_{k=0}^{r_n-1} a_k X^{r_n-k} (X^{s_n} - 1) + \sum_{k=0}^{s_n-1} a_{r_n+k} X^{s_n-k} \in \mathbf{Z}[X],$$

as in Lemma 1. Dividing by $b^{r_n+s_n}$ both sides of (6.1) and letting n tend to infinity along \mathcal{N}_1 , we get that $z_1 = z_3 \alpha$. Hence, if $z_3 = 0$, then $z_1 = 0$, and now relation (6.1) shows that $z_2 = 0$ as well, which is impossible.

Thus, $z_3 \neq 0$ and by multiplying by z_3^{-1} both sides of (6.1), it follows that we may assume that $z_3 = 1$. Since $z_1 = z_3 \alpha$, we get that $z_1 = \alpha$. Inserting this into relation (6.1) and dividing both sides of the resulting relation by $b^{r_n+s_n}$ we get

$$\alpha - \frac{z_2}{b^{s_n}} = \frac{P_n(b)}{b^{r_n+s_n}} = \sum_{k=0}^{s_n+r_n-1} \frac{a_k}{b^k} - \sum_{k=0}^{r_n-1} \frac{a_k}{b^{s_n+k}}.$$

Substituting α for its formula, performing the obvious cancellations and multiplying both sides of the resulting formula by b^{s_n} , we get

$$z_2 = \sum_{k=0}^{r_n-1} \frac{a_k}{b^k} + \sum_{k \geq r_n} \frac{a_{s_n+k}}{b^k}.$$

Since the above relation is true for every n lying in \mathcal{N}_1 , it follows easily that $\alpha = z_2$ because the right hand side of the above relation tends to α when n tends to infinity along \mathcal{N}_1 . Replacing $z_1 = z_2 = \alpha$ in (6.1), we get that

$$\alpha b^{r_n} (b^{s_n} - 1) = P_n(b), \quad (6.2)$$

for every $n \in \mathcal{N}_1$.

Assume first that $\alpha = 0$. We then get that $P_n(b) = 0$. Note that

$$P_n(X) = \sum_{k=0}^{r_n+s_n-1} c_k X^{r_n+s_n-k},$$

where $c_k = a_k$ if $k = 0, \dots, s_n - 1$, and $c_k = a_k - a_{k-s_n}$ if $k = s_n, \dots, r_n + s_n - 1$. Since $a_0 \neq 0$, it follows that $P_n(X)$ is not the zero polynomial. Note that each of its coefficients is either of the form a_i , or of the form $a_j - a_h$ for some appropriate indices i, j and h . Dividing both sides of equation (6.2) by an appropriate power of b to emphasize a constant term (i.e., a term which is not a multiple of b), we get a relation of the form

$$c_{0,n} b^{d_n} + c_{1,n} b^{d_n-1} + \dots + c_{d_n,n} = 0,$$

where $d_n > 0$, the coefficients $c_{j,n}$ are integers, $c_{0,n} = a_0 \neq 0$, $c_{d_n,n} \neq 0$, and the absolute values of all coefficients $c_{i,n}$ does not exceed $|b|^2 - 1$. However, the above relation implies that the polynomial

$$Q_n(X) = c_{0,n}X^{d_n} + c_{1,n}X^{d_n-1} + \dots + c_{d_n,n} \in \mathbf{Z}[X]$$

is a multiple of the minimal polynomial of b over \mathbf{Z} . Since the constant term of the minimal polynomial over \mathbf{Z} of b is $|b|^2 = \bar{b} \cdot b$, and this number must divide the nonzero integer $c_{d_n,n}$ of absolute value $\leq |b|^2 - 1$, we get a contradiction.

From now on, we assume that $\alpha \neq 0$.

Let $n_1 < n_2 < \dots$ be an infinite sequence of positive integers in \mathcal{N}_1 . Thus, relation (6.2) holds for $n = n_i$ with some $i \geq 1$. The sequence $(r_n)_{n \geq 1}$ might be bounded or not. If it is unbounded, we may assume, up to discarding some values for the n_i 's, that $r_{n_{i+1}} > s_{n_i} + r_{n_i}$ for all $i \geq 1$, while if it is bounded, we may assume that r_{n_i} is a constant for all $i \geq 1$. We also assume that $s_{n_{i+1}} > r_{n_i} + s_{n_i}$ for all $i \geq 1$. This is possible because the sequence $(s_n)_{n \geq 1}$ is increasing. From now on, we write $m = n_j$ and $n = n_i$ for some $j > i \geq 1$. Dividing the two relations (6.2) obtained for m and n , we get

$$\frac{b^{r_m}(b^{s_m} - 1)}{b^{r_n}(b^{s_n} - 1)} = \frac{P_m(b)}{P_n(b)},$$

which gives

$$b^{r_m-r_n}(b^{s_m} - 1)P_n(b) - (b^{s_n} - 1)P_m(b) = 0.$$

The expression appearing in the left hand side of the last equation above is a polynomial in b (note that $r_m \geq r_n$). Put $D = (r_m - r_n) + s_m + r_n + s_n = r_m + s_m + r_n$. Note that since $a_0 \neq 0$, it follows that D is the common degree of the polynomials $X^{r_m-r_n}(X^{s_m} - 1)P_n(X)$ and $(X^{s_n} - 1)P_m(X)$. We show that if n is fixed and m is very large, then one of the polynomials

$$Q_{m,n}(X) = X^{r_m-r_n}(X^{s_m} - 1)P_n(X) - (X^{s_n} - 1)P_m(X) \in \mathbf{Z}[X] \quad (6.3)$$

is not the zero polynomial, and then later that $Q_{m,n}(X)$ is of the form $X^\lambda R_{m,n}(X)$, where $R_{m,n}(0)$ is nonzero and of the form $\pm a_i$ or $a_j - a_h$ for some appropriate indices i, j and h , which leads to the same final contradiction as in the case when $\alpha = 0$.

To see that $Q_{m,n}(X)$ is nonzero, note that the coefficient of $X^{D-r_n-s_n-t}$, where $t = 0, 1, \dots, s_m - s_n - r_n$ is simply $a_{s_n+t} - a_t$. If this coefficient is zero for all t in the above range and for arbitrarily large values of m , we then get that the sequence $(a_k)_{k \geq 0}$ is periodic with period s_n , which is a contradiction.

We now look at the last nonzero coefficient of $Q_{m,n}(X)$. Note that if $r_m - r_n > s_n$, then the last nonzero coefficient of $Q_{m,n}(X)$ is $a_{r_m+s_m-1} - a_{r_m-1}$ and it is the coefficient of X . If $a_{r_m+s_m-1} - a_{r_m-1}$ is nonzero, we have already obtained the desired contradiction. If not, we divide both sides of relation (6.3) by X and work with the polynomial $Q_{m,n}(X)/X$. Continuing in this way $r_m - r_n$ times, we may assume that $a_{s_m+r_m-t} = a_{r_m-t}$ for all

$t = 1, \dots, r_m - r_n$. At this point, $Q_{m,n}(X)$ is replaced by polynomial $Q_{m,n}(X)/X^{r_m-r_n}$ given by

$$Q_{m,n}(X)/X^{r_m-r_n} = (X^{s_m} - 1)P_n(X) - (X^{s_n} - 1)P_m(X)/X^{r_m-r_n}. \quad (6.3)$$

Note that if $r_m = r_n$, then the above discussion is unnecessary and the polynomial shown at (6.3) is just $Q_{m,n}(X)$. In what follows, we study the above polynomial. Note that

$$\begin{aligned} S_{m,n}(X) &= (X^{s_m} - 1)P_n(X) \\ &= a_0 X^{s_m+r_n+s_n} + a_1 X^{s_m+r_n+s_n-1} + \dots \\ &\quad + a_{s_n-1} X^{s_m+r_n+1} + (a_{s_n} - a_0) X^{s_m+r_n} + \dots \\ &\quad + (a_{s_n+r_n-1} - a_{r_n-1}) X^{s_m+1} - a_0 X^{r_n+s_n} + \dots \\ &\quad - a_{s_n-1} X^{r_n+1} - (a_{s_n} - a_0) X^{r_n} - \dots - (a_{s_n+r_n-1} - a_{r_n-1}) X. \end{aligned}$$

Thus, with $D_1 = s_m + s_n + r_n$, it follows that

$$S_{m,n}(X) = \sum_{k=0}^{D_1-1} c_k X^{D_1-k},$$

where c_k is given by:

- (i) a_k if $k = 0, \dots, s_n - 1$;
- (ii) $a_k - a_{k-s_n}$ if $k = s_n, \dots, s_n + r_n - 1$;
- (iii) 0 if $k = s_n + r_n, \dots, s_m - 1$;
- (iv) $-a_{k-s_m}$ if $k = s_m, \dots, s_m + s_n - 1$;
- (v) $-(a_{k-s_m} - a_{k-s_m-s_n})$ if $k = s_m + s_n, \dots, s_m + s_n + r_n - 1$.

For the second polynomial appearing in the formula (6.3)

$$T_{m,n}(X) = (X^{s_n} - 1)P_m(X)/X^{r_m-r_n},$$

a similar calculation gives that

$$T_{m,n}(X) = \sum_{k=0}^{D_1-1} d_k X^{D_1-k},$$

where the coefficients d_k are given by:

- (i) a_k if $k = 0, \dots, s_n - 1$;
- (ii) $a_k - a_{k-s_n}$ if $k = s_n, \dots, s_m - 1$;
- (iii)' $a_k - a_{k-s_m} - a_{k-s_n}$ if $k = s_m, \dots, s_m + r_n - 1$;
- (iv)' $-a_{k-s_n}$ if $k = s_m + r_n, \dots, s_m + s_n - 1$;
- (v)' $-(a_{k-s_n} - a_{k-s_n-s_m})$ if $k = s_m + s_n, \dots, s_m + s_n + r_n - 1$,

provided that $s_n > r_n$. When $s_n = r_n$, the same holds except that (iv)' does not exist, while if $s_n < r_n$, then the group (iii)'-(v)' above needs to be replaced by

- (iii)" $a_k - a_{k-s_m} - a_{k-s_n}$ if $k = s_m, \dots, s_m + s_n - 1$;
- (iv)" $a_k - a_{k-s_m} - (a_{k-s_n} - a_{k-s_n-s_m})$ if $k = s_m + s_n, \dots, s_m + r_n - 1$;
- (v)" $-(a_{k-s_n} - a_{k-s_n-s_m})$ if $k = s_m + r_n, \dots, s_m + r_n + s_n - 1$.

It is now easy to obtain the list of coefficients of $Q_{m,n}(X)$. Namely, $c_k - d_k$ equals (independently of the fact that $s_n > r_n$, $s_n = r_n$ or $s_n < r_n$):

- (i) 0 if $k = 0, 1, \dots, s_n + r_n - 1$;
- (ii) $-(a_k - a_{k-s_n})$ if $k = s_n + r_n, \dots, s_m + r_n - 1$;
- (iii) $-(a_{k-s_m} - a_{k-s_n})$ if $k = s_m + r_n, \dots, s_m + s_n + r_n - 1$,

which confirms the claim about the coefficients of $Q_{m,n}(X)$. Thus, Theorem 2 is proved.

□

Proof of Corollary 3. Let \mathbf{a}' be the sequence $a_0, 0, a_1, 0, a_2, 0, \dots$. It is clear that \mathbf{a}' is also stammering. Now the result follows by applying Theorem 2 to the pair $(\mathbf{a}', b^{1/2}i)$. □

Acknowledgments. We are very grateful to the referee for having pointed to us one important mistake and for many valuable suggestions. F. L. thanks Professor Hans Peter Schlickewei for useful advice. This paper was started during a visit of F. L. at the University Strasbourg 1 in July 2005. He warmly thanks the mathematical department for its hospitality. Y. B. and F. L. were supported in part by the joint Project France-Mexico ANUIES-ECOS M01-M02.

References

- [1] B. Adamczewski & Y. Bugeaud, On the complexity of algebraic numbers I. Expansions in integer bases, *Ann. of Math.* **165** (2007), 547–565.
- [2] B. Adamczewski & Y. Bugeaud, Dynamics for β -shifts and Diophantine approximation, *Ergod. Theory Dynam. Syst.*, to appear.
- [3] B. Adamczewski, Y. Bugeaud & F. Luca, Sur la complexité des nombres algébriques, *C. R. Acad. Sci. Paris* **339** (2004), 11–14.
- [4] J.-P. Allouche & J. O. Shallit, *Automatic Sequences: Theory, Applications, Generalizations*, Cambridge University Press, Cambridge, 2003.
- [5] P. Corvaja & U. Zannier, Some new applications of the subspace theorem, *Compositio Math.* **131** (2002), 319–340.
- [6] K. Mahler, Arithmetische Eigenschaften der Lösungen einer Klasse von Funktionalgleichungen, *Math. Ann.* **101** (1929), 342–366.

- [7] K. Mahler, Lectures on Diophantine approximation, Part 1: g -adic numbers and Roth's theorem. University of Notre Dame, Ann Arbor, 1961.
- [8] Ku. Nishioka, *Mahler Functions and Transcendence*, Lecture Notes in Mathematics **1631**, Springer-Verlag, Berlin, 1996.
- [9] H. P. Schlickewei, The p -adic Thue–Siegel–Roth–Schmidt theorem, *Arch. Math. (Basel)* **29** (1977), 267–270.
- [10] W. M. Schmidt, *Diophantine Approximations and Diophantine Equations*, Lecture Notes in Mathematics **1467**, Springer-Verlag, Berlin, 1991.
- [11] M. Waldschmidt, *Diophantine approximation on linear algebraic groups*, Grundlehren der Mathematischen Wissenschaften **326**, Springer-Verlag, Berlin, 2000.

Boris Adamczewski
 CNRS, Université de Lyon, Université Lyon 1
 Institut Camille Jordan
 43 boulevard du 11 novembre 1918
 69622 VILLEURBANNE Cedex (FRANCE)

Yann Bugeaud
 Université Louis Pasteur
 U. F. R. de mathématiques
 7, rue René Descartes
 67084 STRASBOURG Cedex (FRANCE)

`Boris.Adamczewski@math.univ-lyon1.fr`

`bugeaud@math.u-strasbg.fr`

Florian Luca
 Mathematical Institute
 UNAM
 Ap. Postal 61-3 (Xangari), CP 58 089
 Morelia, Michoacán (MEXICO)

`fluca@matmor.unam.mx`