

**Transcendence measures for continued fractions
involving repetitive or symmetric patterns**

Boris ADAMCZEWSKI (Lyon) & Yann BUGEAUD (Strasbourg)

1. Introduction

It was observed long ago (see e.g., [32] or [20], page 62) that Roth's theorem [28] and its p -adic extension established by Ridout [27] can be used to prove the transcendence of real numbers whose expansion in some integer base contains repetitive patterns. This was properly written only in 1997, by Ferenczi and Mauduit [21], who adopted a point of view from combinatorics on words before applying the above mentioned theorems from Diophantine approximation to establish e.g., the transcendence of numbers with a low complexity expansion. Their combinatorial transcendence criterion was subsequently considerably improved in [9] by means of the multidimensional extension of Roth's theorem established by W. M. Schmidt, commonly referred to as the Schmidt Subspace Theorem [29, 30]. As shown in [4], this powerful criterion has many applications and yields among other things the transcendence of irrational real numbers whose expansion in some integer base can be generated by a finite automaton. The latter result was generalized in [7], where we gave transcendence measures for a large class of real numbers shown to be transcendental in [4]. The key ingredient for the proof is then the Quantitative Subspace Theorem [31]. We described in [7] a general method that allows us in principle to get transcendence measures for real numbers that are proved to be transcendental by an application of Roth's or Schmidt's theorem, or one of their extensions.

Besides expansions in integer bases, a classical way to represent a real number is by its continued fraction expansion. There is actually a long tradition in constructing explicit classes of transcendental continued fractions and especially transcendental continued fractions with bounded partial quotients [24, 13, 26, 11]. Again by means of the Schmidt Subspace Theorem, existing results were recently substantially improved in a series of papers [1, 5, 6, 8, 17], providing new classes of transcendental continued fractions. It is the purpose of the present work to show how the Quantitative Subspace Theorem yields transcendence measures for (most of) these numbers, following the approach initiated in [7]. These measures allow us to locate such numbers in the classification of real numbers defined in 1932 by Mahler [23] and recalled below.

For every integer $d \geq 1$ and every real number ξ , we denote by $w_d(\xi)$ the supremum of the exponents w for which

$$0 < |P(\xi)| < H(P)^{-w}$$

has infinitely many solutions in integer polynomials $P(X)$ of degree at most d . Here, $H(P)$ stands for the naïve height of the polynomial $P(X)$, that is, the maximum of the absolute values of its coefficients. Further, we set $w(\xi) = \limsup_{d \rightarrow \infty} (w_d(\xi)/d)$ and, according to Mahler [23],

2000 *Mathematics Subject Classification* : 11J82

B. A. is supported by the ANR through the project "DyCoNum"–JCJC06_134288.

we say that ξ is an

- A -number, if $w(\xi) = 0$;
- S -number, if $0 < w(\xi) < \infty$;
- T -number, if $w(\xi) = \infty$ and $w_d(\xi) < \infty$ for any integer $d \geq 1$;
- U -number, if $w(\xi) = \infty$ and $w_d(\xi) = \infty$ for some integer $d \geq 1$.

An important feature of this classification is that two transcendental real numbers that belong to different classes are algebraically independent. The A -numbers are precisely the algebraic numbers and, in the sense of the Lebesgue measure, almost all numbers are S -numbers. The existence of T -numbers remained an open problem during nearly forty years, until it was confirmed by Schmidt, see Chapter 3 of [16] for references and further results. The set of U -numbers can be further divided in countably many subclasses according to the value of the smallest integer d for which $w_d(\xi)$ is infinite.

Definition 1.1. *Let $\ell \geq 1$ be an integer. A real number ξ is a U_ℓ -number if and only if $w_\ell(\xi)$ is infinite and $w_d(\xi)$ is finite for $d = 1, \dots, \ell - 1$.*

The Liouville numbers are precisely the U_1 -numbers.

To give a flavour of the results proved in the present paper, we quote below a theorem established in 1962 by A. Baker [13].

Theorem (A. Baker). *Consider a quasi-periodic continued fraction*

$$\xi = [a_0, a_1, \dots, a_{n_0-1}, \underbrace{a_{n_0}, \dots, a_{n_0+r_0-1}}_{\lambda_0 \text{ times}}, \underbrace{a_{n_1}, \dots, a_{n_1+r_1-1}}_{\lambda_1 \text{ times}}, \dots],$$

where the notation implies that $n_k = n_{k-1} + \lambda_{k-1}r_{k-1}$ and the λ_k 's indicate the number of times a block of partial quotients is repeated. Suppose that the sequences $(a_n)_{n \geq 0}$ and $(r_n)_{n \geq 0}$ are respectively bounded by A and K . Set

$$L = \limsup_{k \rightarrow +\infty} \lambda_k / \lambda_{k-1}, \quad \ell = \liminf_{k \rightarrow +\infty} \lambda_k / \lambda_{k-1}.$$

If L is infinite and $\ell > 1$, then ξ is a U_2 -number. Furthermore, if L is finite and $\ell > \exp(4A^K)$, then ξ is either an S -number or a T -number.

Baker's theorem shows that the above quasi-periodic continued fractions for which ℓ is sufficiently large cannot include U_d -numbers with $d \geq 3$, that is, there is a gap in the type of transcendental numbers given by them. To the best of our knowledge, this remains up to now the only result in the literature providing a transcendence measure for transcendental numbers defined via their continued fraction expansion as in [24, 13, 26, 11, 1, 5, 6, 8, 17].

The approach of the present paper leads to an improvement of Baker's result. In Theorem 3.2 below, we obtain the same conclusion as in the Theorem above, with the assumption $\ell > \exp(4A^K)$ replaced by the much weaker one $\ell > 1$. In particular, our result does not depend on the values of A or K . The key point in the study of the continued fractions considered by Baker is that some blocks of partial quotients are repeated consecutively an arbitrarily large number of times. Besides these quasi-periodic continued fractions, our method also applies to

continued fractions involving repetitive patterns in a more hidden form, as well as symmetric patterns. An emblematic example is given by the Thue–Morse continued fraction

$$\xi_{\mathfrak{t}} = [1, 2, 2, 1, 2, 1, 1, 2, \dots],$$

whose sequence of partial quotients is the Thue–Morse infinite word on the alphabet $\{1, 2\}$, that is, the n -th partial quotient of ξ is equal to 1 if the sum of the binary digits of n is even and it is equal to 2 otherwise. Although the Thue–Morse sequence is repetitive in the sense of Theorem 2.1 below, it is well-known that it does not contain a finite word repeated consecutively more than twice. The fact that $\xi_{\mathfrak{t}}$ is transcendental was established by M. Queffélec [26] only in 1998. As a consequence of Theorems 2.1 and 4.1, we strengthen her result by proving that $\xi_{\mathfrak{t}}$ is either an S -number or a T -number. This apparently provides the first transcendence measures for real numbers of this shape.

All along this paper, we study continued fractions

$$\xi := [a_0, a_1, \dots, a_\ell, \dots],$$

such that the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded. Here, q_ℓ stands for the denominator of the ℓ -th convergent to ξ . We recall that this condition is in general very easy to check, and is not very restrictive, since it is satisfied by almost all real numbers (with respect to the Lebesgue measure). It is in particular always the case when the sequence $(a_\ell)_{\ell \geq 0}$ is bounded.

Our paper is organized as follows. In Section 2, we state a general result, Theorem 2.1, providing transcendence measures for a large class of continued fractions. These are termed purely stammering continued fractions. Following [1], we then illustrate this result with some applications to various well-known classes of transcendental continued fractions including those arising from Sturmian, morphic and linearly recurrent infinite words. In Section 3, we improve, in the spirit of [5], the theorem of Baker previously mentioned. Transcendence measures for continued fraction involving symmetric patterns are then discussed in Section 4. We gather some auxiliary results in Section 5 and recall in Section 6 our main tool, the Quantitative Subspace Theorem. The proofs of our results are postponed to Sections 7, 8, 9 and 10.

2. Transcendence measures for purely stammering continued fractions

Throughout the present text, we adopt the point of view from combinatorics on words. Let \mathcal{A} be a given set, not necessarily finite. The length of a word W on the alphabet \mathcal{A} , that is, the number of letters composing W , is denoted by $|W|$. For any positive integer ℓ , we write W^ℓ for the word $W \dots W$ (ℓ times repeated concatenation of the word W). We denote by W^∞ the infinite word obtained by concatenation of infinitely many copies of W . For any positive real number x , we denote by W^x the word $W^{\lfloor x \rfloor} W'$, where W' is the prefix of W of length $[(x - \lfloor x \rfloor)|W|]$. Here, and in all what follows, $\lfloor y \rfloor$ and $\lceil y \rceil$ denote, respectively, the integer part and the upper integer part of the real number y . The study of repetitive patterns occurring in finite or infinite words is a classical topic from combinatorics on words. Several exponents have been introduced to measure the presence of such patterns. Among them, the *initial critical exponent* of an infinite word $\mathbf{a} = (a_n)_{n \geq 0}$, denoted by $\text{ice}(\mathbf{a})$, is defined as the supremum of the

real numbers x for which there are arbitrarily long prefixes of \mathbf{a} of the form W^x . Clearly, $\text{ice}(\mathbf{a})$ is always at least equal to 1 and is infinite if \mathbf{a} is a purely periodic sequence. This exponent was introduced in [15] for reasons apparently unrelated to transcendence.

As we will see in the sequel, the initial critical exponent plays an important role in our context. It essentially allows us to distinguish among the large class of continued fractions considered in Theorem 2.1 the U_2 -numbers from the S - or T -numbers. In particular, we show that to decide whether such a continued fraction ξ is a U_2 -number reduces to a purely combinatorial question about the infinite word formed by the partial quotients of ξ .

2.1. Main results

Let $w > 1$ be a real number. We say that a sequence $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ satisfies Condition $(*)_{\widehat{w}}$ if \mathbf{a} is not eventually periodic and if there exists an infinite sequence of finite words $(V_n)_{n \geq 1}$ such that:

- (i) for every integer $n \geq 1$, the word V_n^w is a prefix of the word \mathbf{a} ;
- (ii) the sequence $(|V_n|)_{n \geq 1}$ is (strictly) increasing;
- (iii) the sequence $(|V_{n+1}|/|V_n|)_{n \geq 1}$ is bounded.

We establish the following result.

Theorem 2.1. *Let $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ be a sequence of positive integers satisfying Condition $(*)_{\widehat{w}}$ for some $w > 1$. Let $(p_\ell/q_\ell)_{\ell \geq 1}$ be the sequence of convergents to the real number*

$$\xi := [a_0, a_1, \dots, a_\ell, \dots],$$

and assume that the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded. If $\text{ice}(\mathbf{a})$ is finite, then ξ is either an S -number or a T -number; otherwise, ξ is a U_2 -number. Moreover, if $\text{ice}(\mathbf{a})$ is finite, then there exists a constant c independent of d such that

$$w_d(\xi) \leq \exp(c (\log 3d)^3 (\log \log 3d)^2), \quad (d \geq 1). \quad (2.1)$$

In the case where $w \geq 2$ in Theorem 2.1, we are actually able to derive a better transcendence measure, namely to replace (2.1) by

$$w_d(\xi) \leq \exp(c (\log 3d)^2 (\log \log 3d)), \quad (d \geq 1). \quad (2.2)$$

The fact that the real number ξ is transcendental when the sequence \mathbf{a} only satisfies assumptions (i) and (ii) in the definition of Condition $(*)_{\widehat{w}}$ is the main result of [1]. In the sequel of this section, we point out several consequences of Theorem 2.1 to various classes of continued fractions already considered in the literature and in particular in [1]. The interested reader will find in that paper more explanations motivating the study of these numbers.

2.2. Applications to Sturmian, morphic, and linearly recurrent continued fractions

Let $\alpha > 1$ be an irrational real number and ρ be a positive real number. A sequence $(a_\ell)_{\ell \geq 0}$ on the alphabet $\{a, b\}$ is a *Sturmian sequence* of slope α and intercept ρ if

$$a_\ell = \lfloor (\ell + 1)\alpha + \rho \rfloor - \lfloor \ell\alpha + \rho \rfloor, \quad (\ell \geq 0), \quad (2.3)$$

or if

$$a_\ell = \lceil (\ell + 1)\alpha + \rho \rceil - \lceil \ell\alpha + \rho \rceil, \quad (\ell \geq 0).$$

Such sequences only take two distinct values which are positive integers. For convenience, we will only consider in the sequel Sturmian sequences defined as in (2.3), so that, α and ρ determine for us a unique Sturmian sequence. The real number $\xi_{\alpha, \rho} := [a_0, a_1, a_2, \dots]$ is then termed a Sturmian continued fraction. The main result of [11] is the fact that Sturmian continued fractions are all transcendental. We make here this result more precise.

Theorem 2.2.1. *Let $\alpha > 1$ be an irrational real number and ρ be a positive real number. Let $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ be a Sturmian sequence of slope α and intercept ρ . Then, the real number*

$$\xi_{\alpha, \rho} := [a_0, a_1, \dots, a_\ell, \dots]$$

is a U_2 -number if and only if α has unbounded partial quotients. If α has bounded partial quotients, then $\xi_{\alpha, \rho}$ satisfies the transcendence measure given in (2.1) and is thus either an S -number or a T -number.

It follows from Theorem 2.2.1 that $\xi_{\sqrt{2}, e}$ is either an S -number or a T -number, while $\xi_{e, \sqrt{2}}$ is a U_2 -number. By one of the properties of Mahler's classification recalled in Section 1, these two real numbers are algebraically independent. The proof of Theorem 2.2.1 is postponed to Section 10.

We turn now to an application of Theorem 2.1 to real numbers whose continued fraction expansion can be fixed by a morphism of a free monoid. Given a finite set \mathcal{A} , we denote by \mathcal{A}^* the free monoid generated by \mathcal{A} . The empty word stands for the neutral element of \mathcal{A}^* . A map from \mathcal{A} into \mathcal{A}^* can be extended to an endomorphism of the free monoid \mathcal{A}^* , which itself can be extended by continuity to a map from $\mathcal{A}^* \cup \mathcal{A}^{\mathbb{N}}$ (endowed with its natural topology) into itself. We call *morphism* such a map, so that, a morphism is now allowed to act on infinite words as well. An infinite word \mathbf{a} is a non-trivial fixed point of a morphism σ if there exists a letter a in \mathcal{A} such that $\mathbf{a} = \lim_{n \rightarrow +\infty} \sigma^n(a)$ and if \mathbf{a} is non-eventually periodic. Such an infinite word is said to be recurrent if its first letter occurs at least twice.

Theorem 2.2.2. *Let σ be a morphism defined over a finite subset of the positive integers. Let $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ be an infinite word which is a non-trivial recurrent fixed point of σ and set*

$$\xi := [a_0, a_1, \dots, a_\ell, \dots].$$

Then, ξ is a transcendental number satisfying the transcendence measure given in (2.1) and is thus either an S -number or a T -number.

In Section 8, we explain how Theorem 2.2.2 can be deduced from Theorem 2.1. As noticed in [1] (see the proof of Corollary 2), a non-trivial fixed point of a binary morphism is recurrent. We thus deduce from Theorem 2.2.2 the following result.

Corollary 2.2.3. *Let a and b be two distinct positive integers. Let σ be a binary morphism defined over $\{a, b\}$. Let $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ be a non-trivial fixed point for σ and set*

$$\xi := [a_0, a_1, \dots, a_\ell, \dots].$$

Then, ξ is a transcendental number satisfying the transcendence measure given in (2.1) and is thus either an S -number or a T -number.

We end this section with an application of Theorem 2.1 to linearly recurrent continued fractions. Given a positive integer k , an infinite word \mathbf{a} is said to be k -linearly recurrent if for every positive integer n each word of length n occurring in \mathbf{a} has at least one occurrence in every factor of \mathbf{a} whose length exceeds kn . A sequence is said to be *linearly recurrent* if it is k -linearly recurrent for some positive integer k . A number of famous sequences in combinatorics on words and symbolic dynamics turn out to share this property. The proof of Theorem 5 of [1] shows that an aperiodic linearly recurrent sequence satisfies Condition $(*)_{\widehat{w}}$ for some real number $w > 1$. Furthermore, it is known that the initial critical exponent of a linearly recurrent infinite word is finite (see for instance [18]). It is also clear from the definition that such a word is written over a finite alphabet. We thus deduce from Theorem 2.1 the following result.

Theorem 2.2.4. *Let $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ be an aperiodic linearly recurrent sequence of positive integers. Then, the real number*

$$\xi := [a_0, a_1, \dots, a_\ell, \dots]$$

is a transcendental number satisfying the transcendence measure given in (2.1) and is thus either an S -number or a T -number.

3. Transcendence measures for Maillet–Baker continued fractions

In this section, we are interested in a class of quasi-periodic continued fractions introduced by Maillet [24] and studied by Baker [13, 14]. Baker proved in [13] the transcendence of these continued fractions by using an approach based on Roth’s Theorem for number fields. His results were recently improved in [5] by means of a new application of the Subspace Theorem. Actually, the approach of [5] combined with the method initiated in [7] allows us to derive transcendence measures for such numbers.

We consider a real number $\xi := [a_0, a_1, \dots, a_\ell, \dots]$, where $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ is a non-eventually periodic sequence of positive integers satisfying the following assumption:

There exists an increasing sequence of positive integers $(n_k)_{k \geq 0}$, a sequence of positive integers $(\lambda_k)_{k \geq 0}$, a finite sequence of positive integers b_1, b_2, \dots, b_r , such that for every positive integer k , the sequence \mathbf{a} begins with

$$a_0 a_1 \dots a_{n_k - 1} \underbrace{BB \dots B}_{\lambda_k \text{ times}}, \quad (3.1)$$

where $B := b_1 b_2 \dots b_r$. It is understood here that

$$a_{n_k-r} a_{n_k-r+1} \dots a_{n_k-1} \neq B. \quad (3.2)$$

We now state our main result regarding Maillet–Baker continued fractions.

Theorem 3.1. *Let $\xi := [a_0, a_1, \dots, a_\ell, \dots]$ be defined as above. Let $(p_\ell/q_\ell)_{\ell \geq 1}$ denote the sequence of convergents to ξ and assume that the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded. Assume further that*

$$\liminf_{k \rightarrow \infty} \frac{\lambda_k}{n_k} > 0 \quad (3.3)$$

and

$$\limsup_{k \rightarrow \infty} \frac{n_{k+1}}{n_k} < +\infty. \quad (3.4)$$

Then, ξ is a transcendental number and there exists a constant c independent of d such that

$$w_d(\xi) \leq \exp(c (\log 3d)^3 (\log \log 3d)^2), \quad (3.5)$$

for every positive integer d . In particular, ξ is either an S -number or a T -number.

The proof of this result also relies on the quantitative version of the Subspace Theorem given in Section 6. As an illustration, we obtain that the real number

$$\xi := [1, 2, 2, 1, 2, 1, 1, 1, 2, 1, 1, 1, 1, 1, 1, 2, 1, 1, 1, \dots],$$

whose n -th partial quotient is equal to 2 if n is a power of 2 and to 1 otherwise, is either an S -number or a T -number. Let us also remark that a Maillet–Baker continued fraction satisfying

$$\limsup_{k \rightarrow \infty} \frac{\lambda_k}{n_k} = +\infty$$

corresponds to a U_2 -number for it is extremely well approximated by the quadratic numbers

$$\alpha_k := [a_0, a_1, a_2, \dots, a_{n_k-1}, B^\infty].$$

We note that such a situation cannot occur in Theorem 3.1 because of (3.2) and (3.4).

The proof of Theorem 3.1 can actually be slightly modified to obtain the following improvement of the Theorem of Baker mentioned in the introduction.

Theorem 3.2. *Let us consider the quasi-periodic continued fraction*

$$\xi = [a_0, a_1, \dots, a_{n_0-1}, \underbrace{a_{n_0}, \dots, a_{n_0+r_0-1}}_{\lambda_0 \text{ times}}, \underbrace{a_{n_1}, \dots, a_{n_1+r_1-1}}_{\lambda_1 \text{ times}}, \dots],$$

where the notation implies that $n_{k+1} = n_k + \lambda_k r_k$ and the λ_k 's indicate the number of times a block of partial quotients is repeated. Suppose that the sequences $(a_n)_{n \geq 0}$ and $(r_n)_{n \geq 0}$ are bounded, that $(a_n)_{n \geq 0}$ is not ultimately periodic, and that

$$\liminf_{k \rightarrow \infty} \frac{\lambda_{k+1}}{\lambda_k} > 1.$$

If, moreover,

$$\limsup_{k \rightarrow \infty} \frac{\lambda_{k+1}}{\lambda_k} < +\infty,$$

then the real number ξ is either an S -number or a T -number satisfying (3.5).

4. Transcendence measures for palindromic continued fractions

In this section, we are interested in continued fractions involving some symmetric patterns, whose study was initiated in [6].

We denote the mirror image of a finite word $W := a_1 \dots a_n$ by $\overline{W} := a_n \dots a_1$. Then, a finite word W is a palindrome if $W = \overline{W}$. An infinite word that begins with arbitrarily large palindromes is termed *palindromic*. It was remarked in [6] (Theorem 2.1) that real numbers whose continued fraction expansion is palindromic are either quadratic or transcendental.

Given a palindromic infinite word \mathbf{a} , we denote by $(n_k)_{k \geq 1}$ the increasing sequence of all lengths of palindromic prefixes of \mathbf{a} . Then, the *palindromic density* $d_p(\mathbf{a})$ of \mathbf{a} is defined by

$$d_p(\mathbf{a}) := \left(\limsup_{k \rightarrow +\infty} \frac{n_{k+1}}{n_k} \right)^{-1}.$$

This notion is for instance discussed in [22]. We refine Theorem 2.1 of [6] as follows.

Theorem 4.1. *Let $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ be a non-periodic sequence of positive integers with a positive palindromic density. Let $(p_\ell/q_\ell)_{\ell \geq 1}$ be the sequence of convergents to the real number*

$$\xi := [a_0, a_1, \dots, a_\ell, \dots],$$

and assume that the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded. If $\text{ice}(\mathbf{a})$ is finite, then ξ is either an S -number or a T -number; otherwise, ξ is a U_2 -number. Moreover, if $\text{ice}(\mathbf{a})$ is finite, then there exists a constant c independent of d such that

$$w_d(\xi) \leq \exp(c (\log 3d)^2 (\log \log 3d)), \quad (d \geq 1). \quad (4.1)$$

Actually, it is not difficult to see that a sequence with a positive palindromic density satisfies Condition $(*)_{\widehat{w}}$ for some real number $w > 1$; see for instance the proof of Theorem 4.6 of [3]. Consequently, we could deduce from Theorem 2.1 a result similar to Theorem 4.1 but with a weaker transcendence measure; namely, with (2.1) instead of (4.1). In particular, simultaneous approximations arising from palindromic patterns seem really needed to reach the transcendence measure given by (4.1). The proof of Theorem 4.1 is given in Section 9.

As an illustration of Theorem 4.1, we obtain that the Thue–Morse continued fraction $\xi_{\mathfrak{t}}$ mentioned in the Introduction is either an S -number or a T -number, and it satisfies the transcendence measure given in (4.1).

5. Auxiliary results

For the reader convenience, we first recall some classical results from the theory of continued fractions, whose proofs can be found for example in the book of Perron [25].

Lemma 5.1. *Let $\alpha = [a_0, a_1, \dots, a_n, \dots]$ and $\beta = [b_0, b_1, \dots, b_n, \dots]$ be two real numbers. Assume that there exists a positive integer ℓ such that $a_i = b_i$ for any $i = 0, \dots, \ell$. We then have $|\alpha - \beta| \leq q_\ell^{-2}$, where q_ℓ denotes the denominator of the ℓ -th convergent to α .*

Lemma 5.2. Let $\alpha = [a_0, a_1, \dots, a_\ell, \dots]$ be a real number with convergents $(p_\ell/q_\ell)_{\ell \geq 1}$. Then, for any $\ell \geq 2$, we have

$$\frac{q_{\ell-1}}{q_\ell} = [0, a_\ell, a_{\ell-1}, \dots, a_1].$$

For positive integers a_1, \dots, a_ℓ , denote by $K_\ell(a_1, \dots, a_\ell)$ the denominator of the rational number $[0, a_1, \dots, a_\ell]$. It is commonly called a *continuant*.

Lemma 5.3. For any positive integers a_1, \dots, a_ℓ and any integer k with $1 \leq k \leq \ell - 1$, we have

$$K_\ell(a_1, \dots, a_\ell) = K_\ell(a_\ell, \dots, a_1)$$

and

$$\begin{aligned} K_k(a_1, \dots, a_k) \cdot K_{\ell-k}(a_{k+1}, \dots, a_\ell) &\leq K_\ell(a_1, \dots, a_\ell) \\ &\leq 2 K_k(a_1, \dots, a_k) \cdot K_{\ell-k}(a_{k+1}, \dots, a_\ell). \end{aligned}$$

Lemma 5.4. Let $(a_\ell)_{\ell \geq 1}$ be a sequence of positive integers. For any positive integer ℓ , we have

$$K_\ell(a_1, \dots, a_\ell) \geq 2^{(\ell-1)/2}.$$

We will also make use of the following four auxiliary results.

Lemma 5.5. Let $\alpha = [a_0, a_1, \dots, a_n, \dots]$ and $\beta = [b_0, b_1, \dots, b_n, \dots]$ be two real numbers whose convergents are respectively denoted by $(p_n/q_n)_{n \geq 1}$ and $(r_n/s_n)_{n \geq 1}$. Assume that both sequences $(q_n^{1/n})_{n \geq 1}$ and $(s_n^{1/n})_{n \geq 1}$ are bounded by a real number M . Assume that there exists a positive integer ℓ such that $a_i = b_i$ for any $i = 0, \dots, \ell$ and $a_{\ell+1} \neq b_{\ell+1}$. Then, there exists a positive real number μ , depending only on M , such that

$$|\alpha - \beta| \geq \frac{1}{q_\ell^\mu}.$$

If we assume moreover that the partial quotients of β are bounded by M , then there exists a positive real number c , depending only on M , such that

$$|\alpha - \beta| \geq \frac{c}{q_\ell^2}.$$

Proof. We follow the proof of Lemma 5 from [2]. The constants implicit in \ll and \gg below depend only on M . Set $\alpha' = [a_{\ell+1}, a_{\ell+2}, \dots]$ and $\beta' = [b_{\ell+1}, b_{\ell+2}, \dots]$. Since $a_{\ell+1} \neq b_{\ell+1}$, we have

$$|\alpha' - \beta'| \geq \min \left\{ \frac{1}{a_{\ell+2} + 1}, \frac{1}{b_{\ell+2} + 1} \right\}. \quad (5.1)$$

Furthermore, our assumption on $(q_n^{1/n})_{n \geq 1}$ and $(s_n^{1/n})_{n \geq 1}$ implies that

$$a_{\ell+1}, a_{\ell+2}, b_{\ell+1}, b_{\ell+2} \ll M^\ell,$$

hence

$$\alpha' \ll M^\ell, \quad \beta' \ll M^\ell \quad \text{and} \quad |\alpha' - \beta'| \gg M^{-\ell}. \quad (5.2)$$

The theory of continued fractions gives that

$$\alpha = \frac{p_\ell \alpha' + p_{\ell-1}}{q_\ell \alpha' + q_{\ell-1}} \quad \text{and} \quad \beta = \frac{p_\ell \beta' + p_{\ell-1}}{q_\ell \beta' + q_{\ell-1}},$$

since the first ℓ partial quotients of α and β are the same. We thus obtain that

$$|\alpha - \beta| = \left| \frac{p_\ell \alpha' + p_{\ell-1}}{q_\ell \alpha' + q_{\ell-1}} - \frac{p_\ell \beta' + p_{\ell-1}}{q_\ell \beta' + q_{\ell-1}} \right| = \left| \frac{\alpha' - \beta'}{(q_\ell \alpha' + q_{\ell-1})(q_\ell \beta' + q_{\ell-1})} \right|. \quad (5.3)$$

Together with (5.1) and (5.2), this yields

$$|\alpha - \beta| \gg M^{-3\ell} q_\ell^{-2}. \quad (5.4)$$

We infer from Lemma 5.4 that $q_\ell \geq 2^{(\ell-1)/2}$. Combined with (5.4), this gives the first assertion of our lemma.

For the second assertion, we proceed as above, noticing that

$$|\alpha' - \beta'| \geq 1 - [0, 1, M + 1] = \frac{1}{M + 2} \quad (5.5)$$

and

$$\left| \frac{\alpha' - \beta'}{q_\ell \alpha' + q_{\ell-1}} \right| \geq \frac{1}{4(M + 2)^2 q_\ell}. \quad (5.6)$$

The latter inequality follows from (5.5) if $\alpha' \leq 2\beta'$, since then $\alpha' \leq 2(M + 1)$. If $\alpha' > 2\beta'$, then we simply observe that

$$\left| \frac{\alpha' - \beta'}{q_\ell \alpha' + q_{\ell-1}} \right| \geq \frac{\alpha'}{2(q_\ell \alpha' + q_{\ell-1})} \geq \frac{1}{2(q_\ell + 1)}$$

to get (5.6). Since $\beta' \leq M + 1$, the second assertion of the lemma follows from (5.3) and (5.6). \square

Lemma 5.6. *Let ξ be a real number and $(p_\ell/q_\ell)_{\ell \geq 1}$ be the sequence of convergents to ξ . If the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded, then $w_1(\xi)$ is finite, that is, ξ is not a Liouville number.*

Proof. Let A be a positive real number such that $q_\ell < A^\ell$ for every positive integer ℓ . By Lemma 5.4 we have $q_\ell \geq \sqrt{2}^{\ell-1}$ and thus there exists a positive real number $w > 1$ such that

$$q_{\ell+1} < q_\ell^w, \quad \text{for } \ell \geq 1.$$

By the theory of continued fractions,

$$\left| \xi - \frac{p_\ell}{q_\ell} \right| > \frac{1}{2q_\ell q_{\ell+1}} > \frac{1}{2q_\ell^{w+1}}$$

holds for every positive integer ℓ . Since the best rational approximations to ξ are in the sequence $(p_\ell/q_\ell)_{\ell \geq 1}$, we get $w_1(\xi) \leq w$. Consequently, ξ is not a Liouville number. \square

We further state a classical Liouville inequality which can be found for instance in [16].

Lemma 5.7. *Let β be an algebraic number of degree d and $P(X)$ a non-constant integer polynomial of degree r . Then, if $P(\beta) \neq 0$, we have*

$$|P(\beta)| \geq \frac{1}{(r+1)^{d-1}(d+1)^{r/2}H(P)^{d-1}H(\beta)^r}.$$

If α and β are two non-zero distinct algebraic numbers of degree respectively equal to n and m , then

$$|\alpha - \beta| \geq \frac{2 \max\{2^{-n}(n+1)^{-(m-1)/2}, 2^{-m}(m+1)^{-(n-1)/2}\}}{(n+1)^{m/2}(m+1)^{n/2}H(\alpha)^m H(\beta)^n}.$$

Our last lemma gives a bound for the height of a quotient of algebraic numbers.

Lemma 5.8. *Let α be a real algebraic number of degree $d \geq 2$. Let x_1, x_2, x_3, x_4 be integers with $(x_3, x_4) \neq (0, 0)$ and of absolute values at most A . Then, the height of the algebraic number $\beta = (x_1\alpha + x_2)/(x_3\alpha + x_4)$ is at most equal to $2^{3(d+1)}H(\alpha)^2 A^{2d}$.*

Proof. Using classical inequalities between the naïve height H , the Mahler measure M and the logarithmic Weyl height h , we get that

$$\begin{aligned} H(\beta) &\leq 2^d M(\beta) \leq 2^d e^{dh(\beta)} \leq 2^d e^{dh(x_1\alpha + x_2)} e^{dh(x_3\alpha + x_4)} \\ &\leq 2^d M(x_1\alpha + x_2) M(x_3\alpha + x_4) \\ &\leq 2^d (d+1)H(x_1\alpha + x_2)H(x_3\alpha + x_4) \\ &\leq 2^{3(d+1)}H(\alpha)^2 A^{2d}, \end{aligned}$$

by Lemma A.4 from [16]. □

6. The Quantitative Subspace Theorem

The proofs of our result rely on the quantitative version of the Schmidt Subspace Theorem. The following statement was proved by Evertse [19]. We refer to [19] for the definition of the height of a linear form L , denoted by $\text{Height}(L)$. For our purpose, it is sufficient to stress that it can be bounded from above in terms of the heights of the coefficients of L .

Theorem E. *Let $m \geq 2$, H and d be positive integers. Let L_1, \dots, L_m be linearly independent (over $\overline{\mathbf{Q}}$) linear forms in m variables with real algebraic coefficients. Assume that $\text{Height}(L_i) \leq H$ for $i = 1, \dots, m$ and that the number field generated by all the coefficients of these linear forms has degree at most d . Let ε be a real number with $0 < \varepsilon < 1$. Then, the primitive integer vectors $\mathbf{x} = (x_1, \dots, x_m)$ in \mathbf{Z}^m with $H(\mathbf{x}) \geq H$ and such that*

$$\prod_{i=1}^m |L_i(\mathbf{x})| < |\det(L_1, L_2, \dots, L_m)| \cdot (\max\{|x_1|, \dots, |x_m|\})^{-\varepsilon}$$

lie in at most

$$c_{m,\varepsilon} (\log 4d) (\log \log 4d) \tag{6.1}$$

proper subspaces of \mathbf{Q}^m , where $c_{m,\varepsilon}$ is a constant which only depends on m and ε .

7. Proof of Theorem 2.1

This section is devoted to the proof of Theorem 2.1. We first show how the initial critical exponent allows us to control the approximation by quadratic numbers.

Lemma 7.1. *Let $\mathbf{a} = (a_\ell)_{\ell \geq 1}$ be a non-periodic sequence of positive integers and set*

$$\xi := [0, a_1, \dots, a_\ell, \dots].$$

Denote by $(p_\ell/q_\ell)_{\ell \geq 1}$ the sequence of convergents to ξ . Assume that the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded and that $\text{ice}(\mathbf{a}) = +\infty$. Then ξ is a U_2 -number.

Proof. By assumption, the initial critical exponent of the sequence \mathbf{a} is infinite. Consequently, for every positive integer n , there exists a finite word V_n such that \mathbf{a} begins with the word V_n . Set $\alpha_n = [0, V_n^\infty]$ and denote by s_n the length of the word V_n . Since α_n is a root of the polynomial

$$P_n(X) := q_{s_n-1}X^2 + (q_{s_n} - p_{s_n-1})X - p_{s_n}, \quad (7.1)$$

its height is at most equal to q_{s_n} . The first ns_n partial quotients of ξ and α_n being the same, we infer from Lemmas 5.1 and 5.3 that

$$|\xi - \alpha_n| < q_{ns_n}^{-2} \leq q_{s_n}^{-2n} \leq H(\alpha_n)^{-2n}. \quad (7.2)$$

Since \mathbf{a} is a non-periodic sequence, the set $\{\alpha_n, n \geq 1\}$ is infinite, and (7.2) implies that $w_2(\xi) = +\infty$. By Lemma 5.6, this shows that ξ is a U_2 -number. \square

Next lemma is essentially outlined at the end of [10].

Lemma 7.2. *Let $\mathbf{a} = (a_\ell)_{\ell \geq 1}$ be a sequence of positive integers satisfying Condition $(*)_{\widehat{w}}$ for some $w > 1$. Let $(p_\ell/q_\ell)_{\ell \geq 1}$ be the sequence of convergents to the real number*

$$\xi := [0, a_1, \dots, a_\ell, \dots].$$

Assume that the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded and that $\text{ice}(\mathbf{a})$ is finite. Then, $w_2(\xi)$ is finite.

Proof. Assume that the sequence of finite words $(V_n)_{n \geq 1}$ and the real number w arising from Condition $(*)_{\widehat{w}}$ are fixed. We also assume that $|V_1| \geq 2$. Let $M \geq 3$ be a positive integer such that $\text{ice}(\mathbf{a}) < M$ and $q_\ell < M^\ell$ for $\ell \geq 1$. Denote by l_n the length of the word V_n for $n \geq 1$.

As in the proof of Lemma 7.1, we set $\alpha_n = [0, V_n^\infty]$ and note that α_n is a quadratic number whose height is at most q_{l_n} . Since the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded and the first $\lfloor wl_n \rfloor$ partial quotients of ξ and α_n are the same, we infer from Lemmas 5.1 and 5.3 that

$$|\xi - \alpha_n| < q_{\lfloor wl_n \rfloor}^{-2} < q_{l_n}^{-2-\delta}, \quad (7.3)$$

for some positive real number δ .

On the other hand, since $\text{ice}(\mathbf{a}) < M$, the numbers α_n and ξ cannot have the same first Ml_n partial quotients. The same observation applies for α_n and α_{n+1} . Since the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded by M , we infer from Lemmas 5.3 and 5.5 that there exists a positive real number λ , depending only on M , such that

$$|\xi - \alpha_n| > q_{l_n}^{-\lambda} \quad (7.4)$$

and

$$|\alpha_{n+1} - \alpha_n| > q_{l_n}^{-\lambda}. \quad (7.5)$$

Let N be a positive integer. Extracting if needed a subsequence of the sequence $(V_n)_{n \geq 1}$, we can assume without any loss of generality that there exists a positive integer C_N , depending only on N and on the maximum of the sequence $(|V_{m+1}|/|V_m|)_{m \geq 1}$, such that

$$N|V_n| < |V_{n+1}| < C_N|V_n|, \quad \text{for } n \geq 1. \quad (7.6)$$

Then, Lemma 5.4 implies that

$$q_{l_{n+1}} > 2^{(N/2)|V_n|},$$

while, by assumption,

$$q_{l_n} < M^{|V_n|}.$$

Thus, if N has been chosen large enough, we can guarantee that

$$q_{l_{n+1}} > \max\{(2q_{l_n}^\lambda)^{1/(2+\delta)}, q_{l_n}^2\} \quad (7.7)$$

for every positive integer n . A simple computation shows that it is sufficient to select

$$N > \max\left\{\frac{4 \log M}{\log 2}, \frac{2\lambda \log M}{(2+\delta) \log 2} + \frac{2}{2+\delta}\right\}.$$

On the other hand, we infer from (7.6) that

$$q_{l_{n+1}} < M^{C_N|V_n|},$$

while Lemma 5.4 implies that

$$q_{l_n} > 2^{(|V_n|-1)/2}.$$

This ensures the existence of a positive integer C , depending only on C_N and M , such that

$$q_{l_{n+1}} < q_{l_n}^C, \quad (7.8)$$

for every positive integer n . A simple computation shows that we can take $C = 6C_N \log M$.

By (7.7) and (7.8), we will thus assume in the sequel that

$$\max\{(2q_{l_n}^\lambda)^{1/(2+\delta)}, q_{l_n}^2\} < q_{l_{n+1}} < q_{l_n}^C, \quad (7.9)$$

for every positive integer n .

Combining (7.3) for α_{n+1} with (7.5) and (7.9), we get that $2|\xi - \alpha_{n+1}|$ is at most $|\alpha_n - \alpha_{n+1}|$ and the triangle inequality then yields

$$|\xi - \alpha_n| \geq \frac{1}{2}|\alpha_{n+1} - \alpha_n|. \quad (7.10)$$

Furthermore, setting $c = 1/(4 \cdot 3^{5/2})$, Lemma 5.7 and (7.10) give

$$|\xi - \alpha_n| \geq cH(\alpha_n)^{-2} H(\alpha_{n+1})^{-2} \geq cq_{l_n}^{-2} H(\alpha_{n+1})^{-2}.$$

We then infer from (7.3) and (7.9) that

$$H(\alpha_{n+1}) \geq c^{1/2} q_{l_n}^{\delta/2} > c^{1/2} q_{l_{n+1}}^{\delta/2C}.$$

For n large enough, say $n \geq n_0$, we thus obtain

$$q_{l_n}^{\delta/3C} < H(\alpha_n) \leq q_{l_n}. \quad (7.11)$$

Set $j := \lceil \log(3C/\delta)/\log 2 \rceil$. Using Inequality (7.9), this leads to

$$q_{l_n}^{\delta/3C} < H(\alpha_n) \leq q_{l_n} < q_{l_{n+j}}^{\delta/3C} < H(\alpha_{n+j}) \leq q_{l_{n+j}} < q_{l_n}^{C^j} < H(\alpha_n)^{3C^{j+1}/\delta}.$$

We also deduce from (7.4) and (7.11) that

$$|\xi - \alpha_n| > \frac{1}{H(\alpha_n)^{3\lambda C/\delta}},$$

for every integer $n \geq n_0$.

Set $\sigma = 3\lambda C/\delta$, $\theta = 3C^{j+1}/\delta$ and $\alpha'_n = \alpha_{n_0+(n-1)j}$ for $n \geq 1$. We have shown the existence of a sequence of quadratic irrational numbers $(\alpha'_n)_{n \geq 1}$ satisfying

- (i) $H(\alpha'_n)^{-\sigma} < |\xi - \alpha'_n| < H(\alpha'_n)^{-2-\delta}$;
- (ii) $H(\alpha'_n) < H(\alpha'_{n+1}) < H(\alpha'_n)^\theta$.

At this point, we are ready to prove that $w_2(\xi)$ is finite. Let us consider a quadratic irrational number α whose height satisfies

$$H(\alpha) > (cH(\alpha'_1)^\delta)^{1/2}.$$

If α belongs to the set $\{\alpha'_n, n \geq 1\}$, we simply use (i) to get that

$$|\xi - \alpha| > H(\alpha)^{-\sigma}. \quad (7.12)$$

Let us now assume that α does not belong to the set $\{\alpha'_n, n \geq 1\}$. Then, there exists a unique integer $n > 1$ such that

$$H(\alpha'_{n-1}) < c^{1/\delta} H(\alpha)^{2/\delta} < H(\alpha'_n). \quad (7.13)$$

In particular, Inequality (ii) shows that

$$H(\alpha'_n) < c^{\theta/\delta} H(\alpha)^{2\theta/\delta}. \quad (7.14)$$

Since by assumption α and α'_n denote two distinct quadratic numbers, Lemma 5.7 implies

$$|\alpha - \alpha'_n| \geq 2cH(\alpha)^{-2}H(\alpha'_n)^{-2}. \quad (7.15)$$

We thus infer from (7.13) and (i) that

$$|\alpha - \alpha'_n| > 2|\xi - \alpha'_n|,$$

hence,

$$|\xi - \alpha| \geq \left| |\xi - \alpha'_n| - |\alpha - \alpha'_n| \right| \geq \frac{1}{2}|\alpha - \alpha'_n|.$$

We now infer from (7.14) and (7.15) that

$$|\xi - \alpha| \geq c^{1-2\theta/\delta}H(\alpha)^{-2-4\theta/\delta}.$$

Together with (7.12), the latter inequality shows that $w_2(\xi)$ is finite, concluding the proof. \square

We are now ready to prove Theorem 2.1.

Proof of Theorem 2.1. We keep the notation of Theorem 2.1. Let $\mathbf{a} = (a_\ell)_{\ell \geq 1}$ be a sequence of positive integers satisfying Condition $(*)_{\widehat{w}}$ for some $w > 1$. Let $(p_\ell/q_\ell)_{\ell \geq 1}$ denote the sequence of convergents to the real number

$$\xi := [0, a_1, \dots, a_\ell, \dots].$$

Assume that the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded by some constant A . By Lemma 5.6, $w_1(\xi)$ is finite. Thus, if $\text{ice}(\mathbf{a})$ is infinite, then Lemma 7.1 implies that ξ is a U_2 -number, concluding the proof in that case.

From now on, we assume that $\text{ice}(\mathbf{a})$ is finite. We first observe that $w_2(\xi)$ is finite, in virtue of Lemma 7.2. Consequently, we thus only have to study the approximations to ξ by algebraic numbers of degree at least three. Let $d \geq 3$ be an integer. Let α be an algebraic number of degree d . At several places in the proof below, it is convenient to assume that the height of α is sufficiently large. Let χ be a positive real number such that

$$|\xi - \alpha| < H(\alpha)^{-\chi}.$$

Our aim is to find an upper bound for χ in terms of d . More precisely, we have to prove that

$$\chi < \exp(c(\log 3d)^3 (\log \log 3d)^2) \quad (7.16)$$

for some constant c which does not depend on d .

Extracting if necessary a subsequence of $(V_n)_{n \geq 1}$, we can assume that there exists a constant C such that

$$(iii') \quad 2|V_n| < |V_{n+1}| < C|V_n|, \text{ for every positive integer } n.$$

Set $s_n = |V_n|$ for every positive integer n . Let κ be the unique positive integer such that

$$q_{s_\kappa} \leq H(\alpha) < q_{s_{\kappa+1}}. \quad (7.17)$$

Denote by M_1 the largest integer such that $q_{s_\kappa}^\chi > q_{[ws_\kappa+M_1]}^2$ and observe that

$$|\xi - \alpha| < q_{[ws_\kappa+h]}^{-2} \quad (7.18)$$

for every $1 \leq h \leq M_1$. By definition of M_1 , we have

$$q_{s_\kappa}^\chi < q_{[ws_\kappa+M_1+1]}^2. \quad (7.19)$$

Given a positive integer ℓ , our assumption implies that $q_\ell \leq A^\ell$, while Lemma 5.4 ensures that $q_\ell \geq (\sqrt{2})^{\ell-1}$. We thus get from (7.19) that

$$(\sqrt{2})^{\chi(s_\kappa-1)} \leq A^{2ws_\kappa+M_1+1}$$

and, using (iii'), we obtain

$$(\sqrt{2})^{\chi(s_\kappa-1)} \leq A^{2ws_\kappa C^{M_1+1}}.$$

Consequently, Inequality (7.16) holds if we have

$$M_1 < c_0(\log 3d)^3(\log \log 3d)^2$$

for some constant c_0 which does not depend on d .

We will argue by contradiction. From now on, we assume that

$$M_1 > c_1(\log 3d)^3(\log \log 3d)^2, \quad (7.20)$$

for some constant c_1 , and we will derive a contradiction if c_1 is sufficiently large.

For every integer $n \geq 1$, set

$$\alpha_n = [0, V_n^\infty]$$

and observe as previously that α_n is a root of the quadratic polynomial $P_n(X)$ given by (7.1). Rolle's Theorem and Lemma 5.1 give that

$$|P_n(\xi)| = |P_n(\xi) - P_n(\alpha_n)| \leq 3q_{s_n} |\xi - \alpha_n| < 3q_{s_n} q_{[ws_n]}^{-2}, \quad (n \geq 1), \quad (7.21)$$

since the condition (i) implies that the first $[ws_n]$ partial quotients of ξ and α_n are the same. Furthermore, we infer from the theory of continued fractions that

$$|q_{s_n}\xi - p_{s_n}| < q_{s_n}^{-1} \quad \text{and} \quad |q_{s_n-1}\xi - p_{s_n-1}| < q_{s_n}^{-1}. \quad (7.22)$$

Using again Rolle's Theorem, Inequalities (7.18) and (7.21) imply that

$$|P_{\kappa+h}(\alpha)| < |P_{\kappa+h}(\xi)| + 3q_{s_{\kappa+h}} |\xi - \alpha| < 6q_{s_{\kappa+h}} q_{[ws_{\kappa+h}]}^{-2}, \quad (1 \leq h \leq M_1). \quad (7.23)$$

Inequalities (7.18) and (7.22) ensure that, for $h = 1, \dots, M_1$,

$$|q_{s_{\kappa+h}}\alpha - p_{s_{\kappa+h}}| < 2q_{s_{\kappa+h}}^{-1} \quad \text{and} \quad |q_{s_{\kappa+h}-1}\alpha - p_{s_{\kappa+h}-1}| < 2q_{s_{\kappa+h}}^{-1}. \quad (7.24)$$

We are now going to apply Theorem E to the following system of linear forms:

$$\begin{aligned} L_1(X_1, X_2, X_3, X_4) &= \alpha^2 X_2 + \alpha(X_1 - X_4) - X_3, \\ L_2(X_1, X_2, X_3, X_4) &= \alpha X_1 - X_3, \\ L_3(X_1, X_2, X_3, X_4) &= X_1, \\ L_4(X_1, X_2, X_3, X_4) &= X_2. \end{aligned}$$

Observe first that these linear forms are independent and with algebraic coefficients. Set $\mathcal{N}_1 = \{s_{\kappa+h}, \lfloor M_1/2 \rfloor \leq h \leq M_1\}$ and $\mathcal{P}_1 = \{\mathbf{p}_n = (q_n, q_{n-1}, p_n, p_{n-1}), n \in \mathcal{N}_1\}$. Let n be in \mathcal{N}_1 . Evaluating these linear forms at the integer point \mathbf{p}_n , we infer from Inequalities (7.23) and (7.24) that

$$\prod_{1 \leq i \leq 4} |L_i(\mathbf{p}_n)| < 12q_n^2 q_{\lfloor wn \rfloor}^{-2}.$$

Using again that $\sqrt{2}^{\ell-1} \leq q_\ell \leq A^\ell$ holds for every non-negative integer ℓ , we get that

$$q_{\lfloor wn \rfloor} > q_n^{1+2\eta},$$

for some positive real number η , depending only on w and on A . Thus, if the height of α is large enough, then s_κ is itself sufficiently large to guarantee that

$$\prod_{1 \leq i \leq 4} |L_i(\mathbf{p}_n)| < |\det(L_1, L_2, L_3, L_4)| q_n^{-\eta}.$$

Furthermore, all elements of the set \mathcal{P}_1 are primitive (since p_n and q_n are always relatively prime) and we have $p_n > \text{Height}(L_i)$ for $i = 1, \dots, 4$ and $n \in \mathcal{N}_1$. Moreover, the coefficients of the linear forms L_1, \dots, L_4 generate a number field of degree d . We can thus apply Theorem E with $m = 4$ and $\varepsilon = \eta$. Let T_1 be the upper bound given by (6.1) for the number of exceptional subspaces. Set

$$M_2 := \lfloor M_1/T_1 \rfloor.$$

Since η does not depend on the constant c_1 , Inequality (7.20) ensures the existence of a constant c_2 such that

$$M_2 > c_2(\log 3d)^2(\log \log 3d). \quad (7.25)$$

By the pigeonhole principle, there exists a proper subspace of \mathbf{Q}^4 containing at least M_2 points of \mathcal{P}_1 . Thus, there exist a non-zero integer vector $\mathbf{x} = (x_1, x_2, x_3, x_4)$ and a set of integers $\mathcal{N}_2 \subset \mathcal{N}_1$ with cardinality $r \geq M_2$ such that

$$x_1 q_n + x_2 q_{n-1} + x_3 p_n + x_4 p_{n-1} = 0, \quad (7.26)$$

for every $n \in \mathcal{N}_2$. Let $l_1 < l_2 < \dots < l_r$ denote the elements of \mathcal{N}_2 once ordered.

We are now going to make the following assumption that will be justified thereafter.

Assumption \mathcal{A} : There exist three integers $1 \leq a < b < c \leq \lfloor r/4 \rfloor$ such that the vectors \mathbf{p}_{l_a} , \mathbf{p}_{l_b} and \mathbf{p}_{l_c} are linearly independent.

This makes possible the choice of $\mathbf{x} = \mathbf{p}_{l_a} \wedge \mathbf{p}_{l_b} \wedge \mathbf{p}_{l_c}$ in Equality (7.26). Here, \wedge denotes the vector product in \mathbf{R}^4 . An easy computation shows that with this choice of \mathbf{x} we have

$$h_1 := \max\{|x_1|, |x_2|, |x_3|, |x_4|\} \leq 12q_{l_c}^3. \quad (7.27)$$

Let us remark now that $(x_2, x_4) \neq (0, 0)$ since the vectors (q_{n_1}, p_{n_1}) and (q_{n_2}, p_{n_2}) are not collinear. Furthermore, α being irrational, the real number

$$\beta := -\frac{x_1 + x_3\alpha}{x_2 + x_4\alpha}$$

is well-defined. It then follows from Lemma 5.8 and Inequality (7.27) that

$$H(\beta) \ll q_{l_c}^{6d} H(\alpha)^2, \quad (7.28)$$

where, as below, the constant implied by \ll depends only on d . Since $l_c > s_{\kappa+1}$, we infer from (7.17) that

$$H(\alpha) < q_{l_c}, \quad (7.29)$$

and from (7.17) and (7.28) that

$$H(\beta) < q_{l_c}^{9d}, \quad (7.30)$$

if the height of α has been chosen large enough. Following (7.26), we have

$$\left| \beta - \frac{q_{n-1}}{q_n} \right| = \left| \frac{x_1 + x_3\alpha}{x_2 + x_4\alpha} - \frac{x_1 + x_3p_n/q_n}{x_2 + x_4p_{n-1}/q_{n-1}} \right|,$$

for every $n \in \mathcal{N}_2$. Using (7.22), we thus obtain

$$\left| \beta - \frac{q_{n-1}}{q_n} \right| \ll \frac{h_1^2}{q_n^2 |x_2 + x_4\alpha|^2}.$$

Then, Inequality (7.27) and Liouville's inequality given in Lemma 5.7 yield that

$$\left| \beta - \frac{q_{n-1}}{q_n} \right| \ll h_1^{2d+2} H(\alpha)^2 q_n^{-2} \ll q_{l_c}^{6(d+1)} H(\alpha)^2 q_n^{-2}.$$

Since $d \geq 3$, the height of α can be chosen large enough so that (7.29) implies

$$\left| \beta - \frac{q_{n-1}}{q_n} \right| < q_{l_c}^{7(d+1)} q_n^{-2},$$

for every $n \in \mathcal{N}_2$. Since by assumption $c \leq \lfloor r/4 \rfloor$, $l_{j+1} > 2l_j$ and $r \geq M_2$, we deduce from (7.25) that

$$|q_{l_j}\beta - q_{l_{j-1}}| < q_{l_j}^{-1/2}, \quad (7.31)$$

for every j with $\lfloor r/2 \rfloor \leq j \leq r$. Set $\mathcal{N}'_2 = \{n \in \mathcal{N}_2, n \geq l_{\lfloor r/2 \rfloor}\}$. Denote by M'_2 the cardinality of this set and observe that $M'_2 \geq M_2/2$.

We are now going to apply again Theorem E. Let us consider the following new system of linear forms:

$$L'_1(Y_1, Y_2, Y_3) = \beta Y_1 - Y_2, \quad L'_2(Y_1, Y_2, Y_3) = \alpha Y_1 - Y_3, \quad L'_3(Y_1, Y_2, Y_3) = Y_1.$$

Observe that these linear forms have algebraic coefficients and are linearly independent. Furthermore, $\det(L'_1, L'_2, L'_3) = 1$ and the coefficients of these linear forms generate a field of degree d . Let us also remark that, by Inequality (7.30), we have $\text{Height}(L'_i) < q_n$, for $i = 1, 2, 3$ and every $n \in \mathcal{N}'_2$.

Let n be in \mathcal{N}'_2 . Evaluating these linear forms at the primitive integer point (q_n, q_{n-1}, p_n) , we infer from Inequalities (7.22) and (7.31) that

$$\prod_{1 \leq i \leq 3} |L'_i(q_n, q_{n-1}, p_n)| < q_n^{-1/2}.$$

We can thus apply Theorem E with $\varepsilon = 1/2$ and $m = 3$. Let us denote by T_2 the upper bound given in (6.1) for the number of exceptional subspaces. Set $M_3 := \lfloor T_2/M'_2 \rfloor$. Then, Inequality (7.25) implies the existence of a constant c_3 such that

$$M_3 > c_3 \log d. \quad (7.32)$$

By the pigeonhole principle, there exists a proper subspace of \mathbf{Q}^3 which contains at least M_3 points lying in the set $\mathcal{P}_2 := \{\mathbf{p}_n, n \in \mathcal{N}'_2\}$. Thus, there exist a non-zero integer vector $\mathbf{y} = (y_1, y_2, y_3)$ and a set of integers $\mathcal{N}_3 \subset \mathcal{N}'_2$ with cardinality $s \geq M_3$, and such that

$$y_1 q_n + y_2 q_{n-1} + y_3 p_n = 0, \quad (7.33)$$

for every $n \in \mathcal{N}_3$. Let $m_1 < \dots < m_s$ denote the elements of \mathcal{N}_3 once ordered.

Observe that $\mathbf{p}'_1 = (q_{m_1}, q_{m_1-1}, p_{m_1})$ and $\mathbf{p}'_2 = (q_{m_2}, q_{m_2-1}, p_{m_2})$ are not collinear since $m_2 > m_1 > 1$. This makes possible the choice of $\mathbf{y} = \mathbf{p}'_1 \wedge \mathbf{p}'_2$ in Equality (7.33). Here and in the rest of the proof, \wedge denote the vector product in \mathbf{R}^3 . Then, an easy computation shows that

$$h_2 := \max\{|y_1|, |y_2|, |y_3|\} \leq q_{m_2}^2. \quad (7.34)$$

For every $n \in \mathcal{N}_3$, we can rewrite Equality (7.33) as

$$y_1 + y_2 \beta + y_3 \alpha + y_2 \left(\frac{q_{n-1}}{q_n} - \beta \right) + y_3 \left(\frac{p_n}{q_n} - \alpha \right) = 0.$$

In particular, for $n = m_s$, Inequalities (7.22), (7.31) and (7.34) give

$$|y_1 + y_2 \beta + y_3 \alpha| < 2q_{m_2}^2 q_{m_s}^{-3/2}. \quad (7.35)$$

On the other hand, we have

$$|y_1 + y_2 \beta + y_3 \alpha| = \frac{|y_3 x_4 \alpha^2 + (x_2 y_3 + x_3 y_2 + x_4 y_1) \alpha + (x_1 y_2 + y_1 x_2)|}{|x_2 + x_4 \alpha|}.$$

Thus, if $y_1 + y_2\beta + y_3\alpha \neq 0$, we infer from Liouville's inequality given in Lemma 5.7 that

$$|y_1 + y_2\beta + y_3\alpha| \gg h_1^{-d} h_2^{-d+1} H(\alpha)^{-2}.$$

Using (7.17), (7.27), (7.34) and the fact that $m_2 > l_c > s_{\kappa+1}$, the previous inequality gives

$$|y_1 + y_2\beta + y_3\alpha| \gg q_{m_2}^{-5d}. \quad (7.36)$$

By (7.32), we have $s \geq M_3 > c_3 \log d$ and since by assumption $m_j > 2m_{j-1}$, we can secure that $q_{m_s} > q_{m_2}^{7d}$ and

$$|y_1 + y_2\beta + y_3\alpha| > q_{m_2}^2 q_{m_s}^{-1},$$

if c_3 is chosen large enough. Then, (7.35) contradicts (7.36). Thereby, we get

$$y_1 + y_2\beta + y_3\alpha = 0. \quad (7.37)$$

In order to reach the desired contradiction, we need another equation linking α and β . Actually, it is sufficient to slightly modify the previous system of linear forms and to consider the system:

$$L_1''(Z_1, Z_2, Z_3) = \beta Z_1 - Z_2, \quad L_2''(Z_1, Z_2, Z_3) = \alpha Z_2 - Z_3, \quad L_3''(Z_1, Z_2, Z_3) = Z_1.$$

Again, $\det(L_1'', L_2'', L_3'') = 1$ and the coefficients of L_1'', L_2'', L_3'' generate a number field of degree d . Furthermore, we have $\text{Height}(L_i'') < q_n$ for $i = 1, 2, 3$ and every $n \in \mathcal{N}'_2$. Given an integer $n \in \mathcal{N}'_2$ and evaluating these linear forms at the integer point (q_n, q_{n-1}, p_{n-1}) , we infer from (7.22) and (7.31) that

$$\prod_{1 \leq i \leq 3} |L_i''(q_n, q_{n-1}, p_{n-1})| < q_n^{-1/2}.$$

We can apply Theorem E exactly as before and we will thus omit some details in what follows. The pigeonhole principle ensures the existence of a proper subspace of \mathbf{Q}^3 containing at least M_4 points of \mathcal{P}'_2 . Consequently, there exist a non-zero integer vector $\mathbf{z} = (z_1, z_2, z_3)$ and a set $\mathcal{N}_4 \subset \mathcal{N}'_2$ with cardinality $t \geq M_4$, such that

$$z_1 q_n + z_2 q_{n-1} + z_3 p_{n-1} = 0,$$

for every $n \in \mathcal{N}_4$. Let $n_1 < \dots < n_t$ denote the elements of \mathcal{N}_4 once ordered.

Then, Inequality (7.34) can be replaced by

$$\max\{|z_1|, |z_2|, |z_3|\} \leq q_{n_2}^2,$$

while (7.35) becomes

$$\left| \frac{z_1}{\beta} + z_2 + z_3\alpha \right| < 2q_{n_2}^2 q_{n_t}^{-3/2}.$$

Using Lemma 5.7, we argue as above to show that

$$\frac{z_1}{\beta} + z_2 + z_3\alpha = 0 \quad (7.38)$$

if c_3 is sufficiently large.

Our last step is to show that Equalities (7.37) and (7.38) provide a contradiction. Indeed, since α is irrational, y_2 is not zero and these equalities imply

$$(z_3\alpha + z_2)(y_3\alpha + y_1) = y_2z_1. \quad (7.39)$$

Now, if $y_3z_3 = 0$, then (7.37) and (7.38) imply that β is rational. This gives a contradiction in virtue of (7.30) and (7.31). Consequently, we can assume that $y_3z_3 \neq 0$. Then, we obtain from (7.39) that α is a quadratic irrational. This contradicts the fact that α is an algebraic number of degree at least three. We have proved that the constant c_3 cannot be taken arbitrarily large. Thus, the constant c_1 in (7.20) is bounded, as well. This ends the proof in the case where *Assumption \mathcal{A}* is satisfied.

We turn now to the case where *Assumption \mathcal{A}* is not satisfied. Since some steps are very close from the previous exposition, we just outline the proof.

We come back to the place where *Assumption \mathcal{A}* has been introduced. Set $\mathcal{N}_2'' = \{n \in \mathcal{N}_2, l_{\lfloor r/8 \rfloor} \leq n \leq l_{\lfloor r/4 \rfloor}\}$ and $\mathcal{P}_2'' = \{\mathbf{p}_n, n \in \mathcal{N}_2''\}$. Since *Assumption \mathcal{A}* is not satisfied, all points lying in \mathcal{P}_2'' belong to the subspace generated by \mathbf{p}_{l_1} and \mathbf{p}_{l_2} . Indeed, these vectors are clearly not collinear. In particular, if we set

$$\begin{pmatrix} y'_1 \\ y'_2 \\ y'_3 \end{pmatrix} := \begin{pmatrix} q_{l_1} \\ q_{l_1-1} \\ p_{l_1} \end{pmatrix} \wedge \begin{pmatrix} q_{l_2} \\ q_{l_2-1} \\ p_{l_2} \end{pmatrix},$$

we obtain that (y'_1, y'_2, y'_3) is a non-zero integer vector such that

$$y'_1 q_n + y'_2 q_{n-1} + y'_3 p_n = 0$$

for every integer $n \in \mathcal{N}_2''$. Moreover, an easy computation shows that

$$h'_1 := \max\{|y'_j|, 1 \leq j \leq 3\} < q_{l_2}^2. \quad (7.40)$$

Then, we set

$$\beta' := -\frac{y'_1 + y'_3\alpha}{y'_2}.$$

Note that β' is well-defined since it is easily checked that y'_2 is not zero. Using now (7.40) and Lemma 5.8, we obtain

$$H(\beta') < q_{l_2}^{3d}. \quad (7.41)$$

On the other hand, we can chose the height of α large enough to ensure that

$$|q_n\beta - q_{n-1}| < q_n^{-1/2},$$

for every $n \in \mathcal{N}_2''$. Since \mathcal{N}_2'' is a set with cardinality $M_2'' \geq M_2/8$, we infer from Lemma 5.7 and (7.41) that

$$z'_1 + z'_2\beta + z'_3\alpha = 0. \quad (7.42)$$

Setting

$$\begin{pmatrix} z'_1 \\ z'_2 \\ z'_3 \end{pmatrix} := \begin{pmatrix} q_{l_1} \\ q_{l_1-1} \\ p_{l_1-1} \end{pmatrix} \wedge \begin{pmatrix} q_{l_2} \\ q_{l_2-1} \\ p_{l_2-1} \end{pmatrix},$$

the non-zero integer vector (z'_1, z'_2, z'_3) is such that

$$z'_1 q_n + z'_2 q_{n-1} + z'_3 p_{n-1} = 0,$$

for every $n \in \mathcal{N}_2''$. As previously, this leads to

$$\frac{z'_1}{\beta'} + z'_2 + z'_3 \alpha = 0. \quad (7.43)$$

We then obtain a contradiction using (7.42) and (7.43), exactly as in the case where Assumption \mathcal{A} is satisfied. This concludes the proof of the main part of the theorem.

We now briefly explain why the transcendence measure given by (2.1) can be improved to (2.2) when the sequence \mathbf{a} satisfies Condition $(*)_{\widehat{w}}$ for some $w \geq 2$. In that case the quadratic numbers α_n introduced at the beginning of the present proof are very good approximations of ξ . More precisely, following the proof of Theorem 1 of [1], we obtain that

$$|\xi - \alpha_n| \ll H(\alpha_n)^{-4}.$$

On the other hand, we can assume without loss of generality that the α_n 's are all distinct and that

$$\limsup_{n \rightarrow +\infty} \frac{\log H(\alpha_{n+1})}{\log H(\alpha_n)} < +\infty.$$

The desired transcendence measure then follows from Corollaire 5.3 of [7]. \square

8. Proofs of Theorems 3.1 and 3.2

Proof of Theorem 3.1. We keep the notation of Theorem 3.1 and denote by $A \geq 3$ a strict upper bound for the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$. Without any loss of generality, we assume that $a_0 = 0$.

Let us consider the quadratic real number β whose continued fraction expansion is purely periodic with period $\overline{B} = b_r, b_{r-1}, \dots, b_1$, that is,

$$\beta := [b_r, b_{r-1}, \dots, b_1, b_r, b_{r-1}, \dots, b_1, \dots] = [\overline{B}, \overline{B}, \dots, \overline{B}, \dots].$$

Let $(r_\ell/s_\ell)_{\ell \geq 1}$ denote the sequence of convergents to β . For every positive integer k , set

$$P_k := p_{n_k+r\lambda_k}, \quad Q_k := q_{n_k+r\lambda_k}, \quad P'_k := p_{n_k+r\lambda_k-1}, \quad Q'_k := q_{n_k+r\lambda_k-1} \quad \text{and} \quad S_k := s_{r\lambda_k}.$$

As observed in Section 3, the assumptions (3.2) and (3.4) imply that

$$\limsup_{k \rightarrow +\infty} \frac{\lambda_k}{n_k} < +\infty.$$

Consequently, extracting a subsequence of the sequence $(n_k)_{k \geq 1}$ if necessary, our assumption (3.4) ensures the existence of a constant C_1 such that

$$n_k + r\lambda_k < n_{k+1} < n_{k+1} + r\lambda_{k+1} < C_1(n_k + r\lambda_k)$$

for every positive integer k . Let N be a positive integer. Taking if necessary a subsequence of the sequence $(n_k)_{k \geq 1}$, we can assume without loss of generality that there exists a constant C_N depending on N and C_1 such that

$$N(n_k + r\lambda_k) < n_{k+1} + r\lambda_{k+1} < C_N(n_k + r\lambda_k) \quad (8.1)$$

for every positive integer k . Then, Lemma 5.4 implies that

$$Q_{k+1} > 2^{(N(n_k + r\lambda_k) - 1)/2},$$

while by assumption

$$Q_k < A^{n_k + r\lambda_k}.$$

Thus, if N has been chosen large enough, we obtain that

$$Q_{k+1} > Q_k^2 \quad (8.2)$$

for every positive integer k . A simple computation shows that it is sufficient to choose

$$N > \frac{2 \log A}{\log 2} + 1.$$

On the other hand, we infer from (8.1) that

$$Q_{k+1} < A^{C_N(n_k + r\lambda_k)},$$

while Lemma 5.4 implies that

$$Q_k > 2^{(n_k + r\lambda_k - 1)/2}.$$

This ensures the existence of a positive integer C , depending only on C_N and A , such that

$$Q_{k+1} < Q_k^C, \quad (8.3)$$

for every positive integer k . A simple computation shows that we can take $C = 6C_N \log A$.

By (8.2) and (8.3), we will thus assume in the sequel that

$$Q_k^2 < Q_{k+1} < Q_k^C, \quad (8.4)$$

for every positive integer k .

We also infer from (i) and (ii) that there exists a positive real number ν such that

$$\nu < \lambda_k/n_k < 1/\nu, \quad (8.5)$$

for every positive integer k .

On the one hand, the theory of continued fractions gives

$$|Q_k \xi - P_k| < \frac{1}{Q_k} \quad \text{and} \quad |Q'_k \xi - P'_k| < \frac{1}{Q'_k}, \quad (8.6)$$

while, on the other hand, the assumption made on the sequence \mathbf{a} implies that

$$\frac{P_k}{Q_k} = [0, a_1, \dots, a_{n_k-1}, \underbrace{B, B, \dots, B}_{\lambda_k}].$$

By Lemma 5.2, we obtain

$$\frac{Q'_k}{Q_k} = [0, \underbrace{\overline{B}, \overline{B}, \dots, \overline{B}}_{\lambda_k}, a_{n_k-1}, \dots, a_1]$$

and, following Lemma 5.1, we get that

$$|Q'_k \beta - Q_k| < \frac{Q'_k}{S_k^2}. \quad (8.7)$$

Furthermore, we infer from (3.2) that $a_{n_k-1} \dots a_{n_k-r} \neq \overline{B}$. Thus, Lemmas 5.3 and 5.5 imply that

$$\left| \beta - \frac{Q_k}{Q'_k} \right| > \frac{c_\beta}{S_k^2}, \quad (8.8)$$

where c_β is a positive constant depending only on β .

Since the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded, we infer from Lemma 5.6 that $w_1(\xi)$ is finite. In the sequel, we thus only have to deal with the approximations of ξ by algebraic numbers of degree at least equal to two. Let $d \geq 2$ be an integer. Let α be an algebraic number of degree d , the height of which will be chosen large enough in the sequel. Let χ be a positive real number such that

$$|\xi - \alpha| < H(\alpha)^{-\chi}.$$

Our aim is to find an upper bound for χ as a function of d . More precisely, we have to prove that

$$\chi < \exp(c (\log 3d)^3 (\log \log 3d)^2) \quad (8.9)$$

for some constant c which does not depend on d .

Let us denote by k_0 the unique positive integer such that

$$Q_{k_0} \leq H(\alpha) < Q_{k_0+1}, \quad (8.10)$$

and by M_1 the largest integer such that $Q_{k_0}^\chi > Q_{k_0+M_1}^2$. Consequently,

$$|\xi - \alpha| < Q_{k_0+h}^{-2} \quad (8.11)$$

holds for every integer $h = 1, \dots, M_1$. If

$$M_1 < c_0(\log 3d)^3(\log \log 3d)^2$$

holds for some constant c_0 which does not depend on d , then Inequality (8.9) also holds, as shown by an easy computation using (8.4), (8.10) and (8.11).

We will argue by contradiction. From now on, we assume that

$$M_1 > c_1(\log 3d)^3(\log \log 3d)^2, \quad (8.12)$$

for some constant c_1 , and we will derive a contradiction if c_1 is sufficiently large.

It follows from Inequalities (8.6) and (8.11) that, for every integer $h = 1, \dots, M_1$,

$$|Q_{k_0+h}\alpha - P_{k_0+h}| < \frac{2}{Q_{k_0+h}} \quad \text{and} \quad |Q'_{k_0+h}\alpha - P'_{k_0+h}| < \frac{2}{Q'_{k_0+h}}. \quad (8.13)$$

We are now going to apply Theorem E to the following system of linear forms:

$$\begin{aligned} L_1(X_1, X_2, X_3, X_4) &= \alpha X_1 - X_3, \\ L_2(X_1, X_2, X_3, X_4) &= \alpha X_2 - X_4, \\ L_3(X_1, X_2, X_3, X_4) &= \beta X_1 - X_2, \\ L_4(X_1, X_2, X_3, X_4) &= X_1. \end{aligned}$$

Observe that these linear forms are linearly independent over $\overline{\mathbf{Q}}$ and all have real algebraic coefficients. Set $\mathcal{N}_1 = \{k_0 + h, \lfloor M_1/2 \rfloor \leq h \leq M_1\}$ and $\mathcal{P}_1 = \{\mathbf{p}_k = (Q_k, Q'_k, P_k, P'_k), k \in \mathcal{N}_1\}$. Let $k \in \mathcal{N}_1$. Evaluating these linear forms at the integer point \mathbf{p}_k , Inequalities (8.7) and (8.13) give

$$\prod_{1 \leq i \leq 4} |L_i(\mathbf{p}_k)| < \frac{4}{S_k^2}. \quad (8.14)$$

By assumption, $Q_k \leq A^{n_k+r\lambda_k}$, while Lemma 5.4 implies that $S_k \geq (\sqrt{2})^{r\lambda_k-1}$. We then infer from Inequality (8.5) that

$$S_k \geq Q_k^\eta, \quad (8.15)$$

for some positive real number η , depending only on r , ν and A . If the height of α is large enough, we get from (8.14) and (8.15) that

$$\prod_{1 \leq i \leq 4} |L_i(\mathbf{p}_k)| < |\det(L_1, L_2, L_3, L_4)| Q_k^{-\eta}.$$

Since P_k and Q_k are relatively prime, every element in \mathcal{P}_1 is a primitive vector. Furthermore, the degree of the number field generated by the coefficients of our four linear forms is at most $2d$ and $Q_k > \text{Height}(L_i)$ for $i = 1, \dots, 4$ and $k \in \mathcal{N}_1$. We can thus apply Theorem E with $m = 4$ and $\varepsilon = \eta$. Let T_1 be the upper bound given in (6.1) for the number of exceptional subspaces. Set

$$M_2 := \lfloor M_1/T_1 \rfloor.$$

Since η does not depend on the choice of the constant c_1 , Inequality (8.12) ensures the existence of a constant c_2 which does not depend on d and such that

$$M_2 > c_2(\log 3d)^2(\log \log 3d). \quad (8.16)$$

arbitrarily large in the sequel. By the pigeonhole principle, there exists a proper subspace of \mathbf{Q}^4 containing at least M_2 points of \mathcal{P}_1 . That is, there exist a non-zero integer vector $\mathbf{x} = (x_1, x_2, x_3, x_4)$ and a set of integers $\mathcal{N}_2 \subset \mathcal{N}_1$ with cardinality $s \geq M_2$ such that

$$x_1Q_k + x_2Q'_k + x_3P_k + x_4P'_k = 0, \quad (8.17)$$

for every $k \in \mathcal{N}_2$. Let $l_1 < \dots < l_s$ denote the elements of \mathcal{N}_2 once ordered.

We are now going to make the following assumption that will be justified thereafter.

Assumption \mathcal{A} : we assume that there exist three integers $1 \leq a < b < c \leq \lfloor s/4 \rfloor$ such that the vectors \mathbf{p}_{l_a} , \mathbf{p}_{l_b} and \mathbf{p}_{l_c} are linearly independent.

This makes possible the choice of $\mathbf{x} = \mathbf{p}_{l_a} \wedge \mathbf{p}_{l_b} \wedge \mathbf{p}_{l_c}$ in Equality (8.17). Here, \wedge denotes the vector product in \mathbf{R}^4 . As shown by an easy computation, this choice of \mathbf{x} guarantee that

$$h_1 := \max\{|x_1|, |x_2|, |x_3|, |x_4|\} \leq 12Q_{l_c}^3. \quad (8.18)$$

It thus follows from Inequality (8.17) that

$$|x_4\alpha Q'_k + x_1Q_k + x_2Q'_k + x_3P_k| = |x_4(Q'_k\alpha - P'_k)| < \frac{|x_4|}{Q_k} < \frac{12Q_{l_c}^3}{Q_k}, \quad (8.19)$$

for every $k \in \mathcal{N}_2$.

Our next step consists in applying again Theorem E to the following new system of linear forms:

$$\begin{aligned} L'_1(X_1, X_2, X_3) &= \alpha X_1 - X_3, \\ L'_2(X_1, X_2, X_3) &= x_1X_1 + (x_4\alpha + x_2)X_2 + x_3X_3, \\ L'_3(X_1, X_2, X_3) &= X_1. \end{aligned}$$

We first observe that these linear forms have algebraic coefficients. We also deduce from Equality (8.17) that $(x_2, x_4) \neq (0, 0)$. Consequently, $x_4\alpha + x_2 \neq 0$, since α is irrational. We then deduce that our linear forms are linearly independent. Furthermore, the number field generated by all the coefficients of these linear forms has degree d , and we infer from Lemma 5.7, (8.18) and (8.19) that

$$|\det(L'_1, L'_2, L'_3)| = |x_4\alpha + x_2| > Q_{l_c}^{-3d}, \quad (8.20)$$

provided that $H(\alpha)$ is large enough.

Since by assumption $c \leq \lfloor s/4 \rfloor$ and $s \geq M_2$, we infer from (8.4) and (8.16) that the constant c_2 can be chosen large enough so that

$$Q_k^{1/2} > 12 Q_{l_c}^{3d+3}, \quad (8.21)$$

for every $\lfloor s/2 \rfloor \leq k \leq s$. For the same reason, we can assume that $H(L'_i) < Q_k$, for $i = 1, 2, 3$ and every $\lfloor s/2 \rfloor \leq k \leq s$. Set $\mathcal{N}'_2 = \{n \in \mathcal{N}_2, n \geq \lfloor s/2 \rfloor\}$. Let us denote by M'_2 the cardinality of this set and observe that

$$M'_2 \geq M_2/2. \quad (8.22)$$

Given an integer $k \in \mathcal{N}'_2$ and evaluating these linear forms at the primitive integer point $\mathbf{p}'_k := (Q_k, Q'_k, P_k)$, we infer from Inequalities (8.19), (8.20) and (8.21) that

$$\prod_{1 \leq i \leq 3} |L'_i(Q_k, Q'_k, P_k)| < |\det(L'_1, L'_2, L'_3)| Q_k^{-1/2}.$$

Furthermore, it follows from Lemma 5.7 that P_k is larger than $\text{Height}(L'_i)$ for $i = 1, 2, 3$. We can thus apply Theorem E with $\varepsilon = 1/2$ and $m = 3$. Let us denote by T_2 the upper bound given in (6.1) for the number of exceptional subspaces. Set $M_3 := \lfloor T_2/M'_2 \rfloor$. Then, Inequalities (8.16) and (8.22) imply the existence of a constant c_3 such that

$$M_3 > c_3 \log d. \quad (8.23)$$

By the pigeonhole principle, there exists a proper subspace of \mathbf{Q}^3 which contains at least M_3 points lying in the set $\mathcal{P}_2 := \{\mathbf{p}'_k = (Q_k, Q'_k, P_k), k \in \mathcal{N}'_2\}$. There thus exist a non-zero integer vector $\mathbf{y} = (y_1, y_2, y_3)$ and a set of integers $\mathcal{N}_3 \subset \mathcal{N}'_2$ of cardinality $t \geq M_3$, and such that

$$y_1 Q_k + y_2 Q'_k + y_3 P_k = 0, \quad (8.24)$$

for every $k \in \mathcal{N}_3$. Let $1 \leq m_1 < \dots < m_t$ denote the elements of \mathcal{N}_3 once ordered.

Observe that \mathbf{p}'_1 and \mathbf{p}'_2 are not collinear. This makes possible the choice of $\mathbf{y} = \mathbf{p}'_1 \wedge \mathbf{p}'_2$ in Equality (8.24). Here and in the rest of the proof, \wedge denote the vector product in \mathbf{R}^3 . Then, an easy computation shows that

$$h_2 := \max\{|y_1|, |y_2|, |y_3|\} \leq Q_{m_2}^2. \quad (8.25)$$

For every $k \in \mathcal{N}_3$, we can rewrite Equality (8.24) as

$$y_1 + y_2 \beta + y_3 \alpha + y_2 \left(\frac{Q'_k}{Q_k} - \beta \right) + y_3 \left(\frac{P_k}{Q_k} - \alpha \right) = 0. \quad (8.26)$$

Choosing $k = m_t$, we can argue as in the proof of Theorem 2.1 and use (8.23) to deduce from the Liouville inequality given in Lemma 5.7 that

$$y_1 + y_2 \beta + y_3 \alpha = 0. \quad (8.27)$$

As a particular instance of Equality (8.26), we thus obtain

$$y_2 \left(\frac{Q'_{m_t}}{Q_{m_t}} - \beta \right) = y_3 \left(\frac{P_{m_t}}{Q_{m_t}} - \alpha \right).$$

It follows from Lemma 5.3 and (8.5) that there exists a real number $C_1 > 1$, depending only on ν , such that

$$Q_{m_t} > C_1^{m_t} S_{m_t}.$$

Consequently, on the one hand, we have

$$\left| y_3 \left(\frac{P_{m_t}}{Q_{m_t}} - \alpha \right) \right| < \frac{|y_3|}{Q_{m_t}^2} < \frac{|y_3|}{C_1^{2m_t} S_{m_t}^2} < \frac{1}{C_1^{m_t}} \cdot \frac{1}{S_{m_t}^2}, \quad (8.28)$$

by (8.25), if c_3 is sufficiently large. On the other hand, we infer from (8.27) that $y_2 \neq 0$ (since α is irrational), and Inequality (8.8) shows that

$$\left| y_2 \left(\frac{Q'_{m_t}}{Q_{m_t}} - \beta \right) \right| > \frac{c\beta}{S_{m_t}^2}. \quad (8.29)$$

Inequalities (8.28) and (8.29) provide us with a contradiction as soon as c_3 is chosen large enough. This ends the proof in the case where *Assumption* \mathcal{A} is satisfied.

We turn now to outline the case where this assumption is not satisfied. We come back to the place where *Assumption* \mathcal{A} has been introduced. Set $\mathcal{N}_2'' = \{n \in \mathcal{N}_2, l_{\lfloor r/8 \rfloor} \leq n \leq l_{\lfloor r/4 \rfloor}\}$ and $\mathcal{P}_2'' = \{\mathbf{p}_n, n \in \mathcal{N}_2''\}$. Since *Assumption* \mathcal{A} is not satisfied, all points lying in \mathcal{P}_2'' belong to the vectorial plan generated by \mathbf{p}_{l_1} and \mathbf{p}_{l_2} . Indeed, these vectors are clearly not collinear. In particular, if we set

$$\begin{pmatrix} y'_1 \\ y'_2 \\ y'_3 \end{pmatrix} := \begin{pmatrix} q_{l_1} \\ q_{l_1-1} \\ p_{l_1} \end{pmatrix} \wedge \begin{pmatrix} q_{l_2} \\ q_{l_2-1} \\ p_{l_2} \end{pmatrix},$$

we obtain that (y'_1, y'_2, y'_3) is a non-zero integer vector such that

$$y'_1 Q_k + y'_2 Q'_k + y'_3 P_k = 0$$

for every integer $k \in \mathcal{N}_2''$. Moreover, an easy computation shows that

$$h'_1 := \max\{|y'_j|, 1 \leq j \leq 3\} < Q_{l_2}^2.$$

We are now in the same situation as in (8.25). We can thus argue exactly as previously to conclude the proof of the theorem. \square

Proof of Theorem 3.2. The proof of Theorem 3.2 follows essentially the same lines as that of Theorem 3.1, with, however, some modification that we point out below.

Since the sequences $(a_n)_{n \geq 0}$ and $(r_n)_{n \geq 0}$ are bounded, there exists a finite set $\mathcal{B} = \{B_1, \dots, B_m\}$ of finite words such that all the finite words $a_{n_k} a_{n_k+1} \dots a_{n_k+r_k-1}$ belong to the set \mathcal{B} .

For $j = 1, \dots, m$, set

$$\beta_j := [\overline{B_j}, \overline{B_j}, \dots, \overline{B_j}, \dots].$$

Let $(r_\ell^{(j)}/s_\ell^{(j)})_{\ell \geq 1}$ denote the sequence of convergents to β_j . For every positive integer k , set

$$P_k := p_{n_k+r_k \lambda_k}, \quad Q_k := q_{n_k+r_k \lambda_k}, \quad P'_k := p_{n_k+r_k \lambda_k-1}, \quad Q'_k := q_{n_k+r_k \lambda_k-1} \quad \text{and} \quad S_k := s_{r_k \lambda_k}^{(j)},$$

where the index j is such that $a_{n_k} a_{n_k+1} \dots a_{n_k+r_k-1} = B_j$.

Then, we follow the proof of Theorem 3.1 and we consider, for $j = 1, \dots, m$, the following system of linear forms:

$$\begin{aligned} L_1^{(j)}(X_1, X_2, X_3, X_4) &= \alpha X_1 - X_3, \\ L_2^{(j)}(X_1, X_2, X_3, X_4) &= \alpha X_2 - X_4, \\ L_3^{(j)}(X_1, X_2, X_3, X_4) &= \beta_j X_1 - X_2, \\ L_4^{(j)}(X_1, X_2, X_3, X_4) &= X_1. \end{aligned}$$

Similar arguments as above show that, for k in \mathcal{N}_1 , there exist an absolute positive number η and an index j in $\{1, \dots, m\}$ such that

$$\prod_{1 \leq i \leq 4} |L_i^{(j)}(\mathbf{p}_k)| < |\det(L_1^{(j)}, L_2^{(j)}, L_3^{(j)}, L_4^{(j)})| Q_k^{-\eta}.$$

By the pigeonhole principle, there exist a non-zero integer vector $\mathbf{z} = (z_1, z_2, z_3, z_4)$, an index j in $\{1, \dots, m\}$, and a set of integers $\mathcal{N}_3 \subset \mathcal{N}_1$ whose cardinality is at least $\lfloor M_2/m \rfloor$ (with M_2 defined in (8.16)) and such that

$$z_1 Q_k + z_2 Q'_k + z_3 P_k + z_4 P'_k = 0$$

and $a_{n_k} a_{n_k+1} \dots a_{n_k+r_k-1} = B_j$ for every $k \in \mathcal{N}_3$.

From now on, we simply follow the proof of Theorem 3.1, with β replaced by β_j in (8.26) and below. This proves Theorem 3.2. \square

9. Proof of Theorem 4.1

In this section, we explain how Theorem 4.1 can be deduced from Theorem 2.1 and Théorème 4.1 from [7].

Proof of Theorem 4.1. Let $\mathbf{a} = (a_\ell)_{\ell \geq 1}$ be a sequence of positive integers satisfying the assumptions of Theorem 4.1. In particular, \mathbf{a} has a positive palindromic density and we denote by $(n_k)_{k \geq 1}$ the increasing sequence of all lengths of palindromic prefixes of \mathbf{a} . Set $\xi = [0, a_1, \dots, a_\ell, \dots]$ and let us denote as usual by p_ℓ/q_ℓ the ℓ -th convergent to ξ .

As already mentioned in Section 4, the fact that \mathbf{a} satisfies Condition $(*)_{\widehat{w}}$ for some $w > 1$ can be established following the proof of Theorem 4.6 of [3]. It thus follows from Theorem 2.1 that ξ is a U_2 -number if $\text{ice}(\mathbf{a})$ is infinite, while it is either an S -number or a T -number otherwise.

From now on, we assume that $\text{ice}(\mathbf{a})$ is finite. Since the word $a_1 \dots a_{n_k}$ is a palindrome, Lemma 5.2 implies that $p_{n_k} = q_{n_k-1}$. Recalling that

$$\left| \xi - \frac{p_{n_k}}{q_{n_k}} \right| < \frac{1}{q_{n_k}^2} \quad \text{and} \quad \left| \xi - \frac{p_{n_k-1}}{q_{n_k-1}} \right| < \frac{1}{q_{n_k-1}^2},$$

we infer from $p_{n_k} = q_{n_k-1}$, $0 < \xi < 1$, $a_1 = a_{n_k}$, $|p_{n_k} q_{n_k-1} - p_{n_k-1} q_{n_k}| = 1$ and from $q_{n_k} \leq (a_{n_k} + 1) q_{n_k-1}$ that

$$\begin{aligned} \left| \xi^2 - \frac{p_{n_k-1}}{q_{n_k}} \right| &= \left| \xi^2 - \frac{p_{n_k-1}}{q_{n_k-1}} \cdot \frac{p_{n_k}}{q_{n_k}} \right| \leq \left| \xi + \frac{p_{n_k-1}}{q_{n_k-1}} \right| \cdot \left| \xi - \frac{p_{n_k}}{q_{n_k}} \right| + \frac{1}{q_{n_k} q_{n_k-1}} \\ &\leq 2 \left| \xi - \frac{p_{n_k}}{q_{n_k}} \right| + \frac{1}{q_{n_k} q_{n_k-1}} < \frac{a_1 + 3}{q_{n_k}^2}. \end{aligned}$$

Consequently, there exists an increasing sequence of positive integers $(q_{n_k})_{k \geq 1}$ such that

$$\|q_{n_k} \xi\| \cdot \|q_{n_k} \xi^2\| \ll \frac{1}{q_{n_k}^2},$$

where $\|\cdot\|$ denotes the distance to the nearest integer. Furthermore, since \mathbf{a} has a positive palindromic density and the sequence $(q_\ell^{1/\ell})_{\ell \geq 1}$ is bounded, Lemma 5.4 ensures that

$$\limsup_{k \rightarrow +\infty} \frac{\log q_{n_{k+1}}}{\log q_{n_k}} < +\infty.$$

The transcendence measure given in (4.1) is then obtained as a consequence of Théorème 4.1 from [7]. This concludes the proof. \square

10. Proofs of Theorems 2.2.1 and 2.2.2

In this section, we explain how Theorems 2.2.1 and 2.2.2 can be deduced from Theorem 2.1.

Proof of Theorem 2.2.1. Let $\mathbf{a} = (a_\ell)_{\ell \geq 0}$ be a Sturmian sequence of slope α and let a and b be the two integer values taken by \mathbf{a} .

If α has bounded partial quotients, it is proved in [18] that \mathbf{a} is linearly recurrent. We thus can apply Theorem 2.2.4 to conclude the proof in that case.

If α has unbounded partial quotients, the key auxiliary ingredient is Proposition 12.1 from [7]. It asserts that, for every integer m , there exist finite words U_m, V_m and a positive real number s_m such that $U_m V_m^{s_m}$ is a prefix of \mathbf{a} and

$$|U_m V_m^{s_m}| / |U_m V_m| \geq m. \quad (10.1)$$

In this case, the quadratic real number ξ_m whose continued fraction expansion is given by the ultimately periodic infinite word $U_m V_m^\infty$ is a very good approximation to ξ . More precisely, since the height of ξ_m is at most $(a+b)^{|U_m V_m|}$ and, by Lemmas 5.1 and 5.4,

$$|\xi - \xi_m| \leq 2^{-|U_m + s_m V_m| + 1},$$

we deduce from (10.1) that there exists a positive real number C independent of m and such that

$$|\xi - \xi_m| \leq H(\xi_m)^{-Cm}.$$

Consequently, if α has unbounded partial quotients, then ξ is a U_2 -number. \square

Proof of Theorem 2.2.2. Let \mathbf{a} be a non-trivial recurrent fixed point of a morphism σ defined over a finite set \mathcal{A} . In view of Theorem 2.1, it is sufficient to prove that

- the sequence \mathbf{a} satisfies Condition $(*)_{\widehat{w}}$ for some real number $w > 1$;
- $\text{ice}(\mathbf{a})$ is finite.

For the first point, we refer the reader to the proof of Theorem 3 of [1]. It thus remains to prove that $\text{ice}(\mathbf{a})$ is finite. Let us denote by $|\sigma|$ the length of the morphism σ defined by

$$|\sigma| := \max\{|\sigma(a)|, a \in \mathcal{A}\}.$$

We will actually prove that $\text{ice}(\mathbf{a}) \leq 2|\sigma|$.

We assume that this inequality does not hold and aim at deriving a contradiction. Setting $N = 2|\sigma|$, there exists a non-empty word U such that U^N is a prefix of \mathbf{a} . Without loss of generality we can assume that U has been chosen with minimal length. Since \mathbf{a} is fixed by σ , the word $\sigma(U)$ is a prefix of \mathbf{a} . We observe that N is large enough to ensure that $\sigma(U)$ is also a prefix of the word U^N . There thus exist a non-negative integer r and a (possibly empty) prefix U_1 of U such that $|U_1| < |U|$ and

$$\sigma(U) = U^r U_1. \quad (10.2)$$

If U_1 is the empty word, then we deduce that $\mathbf{a} = U^\infty$, providing a contradiction with the fact that \mathbf{a} is a non-eventually periodic sequence.

From now on, we assume that U_1 is not the empty word. Since U^2 is a prefix of \mathbf{a} , the word $\sigma(U^2)$ is also a prefix of \mathbf{a} , and we observe that N is large enough to guarantee that $\sigma(U^2)$ is a prefix of U^N . From (10.2), we infer that $U^r U_1 U^r U_1$ is a prefix of U^N . Let U_2 denote the prefix of U of length $|U| - |U_1|$ and U_3 denote the suffix of U of length $|U_1|$, so that $U = U_2 U_3$. Since $U^r U_1 U^r U_1$ is a prefix of U^N , we deduce that $U = U_1 U_2 = U_3 U_2$. Consequently, the two words U_2 and U_3 commute. Since U_1 and U_2 are two non-empty words, this implies that there exist a word V and an integer $m \geq 2$ such that $U = V^m$. In particular, the sequence \mathbf{a} begins with the word V^N . This contradicts the fact that U has been chosen with minimal length, concluding the proof. \square

Acknowledgements. We are very much indebted to the referee, who pointed out several obscure points in a first version of the present text.

References

- [1] B. Adamczewski and Y. Bugeaud, *On the complexity of algebraic numbers II. Continued fractions*. Acta Math. 195 (2005), 1–20.
- [2] B. Adamczewski and Y. Bugeaud, *On the Littlewood conjecture in simultaneous Diophantine approximation*. J. London Math. Soc. 73 (2006), 355–366.
- [3] B. Adamczewski and Y. Bugeaud, *Real and p -adic expansions involving symmetric patterns*, Int. Math. Res. Not. (2006), Art. ID 75968, 17 pp.
- [4] B. Adamczewski and Y. Bugeaud, *On the complexity of algebraic numbers I. Expansions in integer bases*. Ann. of Math. 165 (2007), 547–565.
- [5] B. Adamczewski and Y. Bugeaud, *On the Maillat–Baker continued fractions*, J. reine angew. Math. 606 (2007), 105–121.

- [6] B. Adamczewski and Y. Bugeaud, *Palindromic continued fractions*, Ann. Inst. Fourier (Grenoble) 57 (2007), 1557–1574.
- [7] B. Adamczewski et Y. Bugeaud, *Mesures de transcendance et aspects quantitatifs de la méthode de Thue–Siegel–Roth–Schmidt*. Preprint.
- [8] B. Adamczewski, Y. Bugeaud and J. L. Davison, *Continued fractions and transcendental numbers*, Ann. Inst. Fourier (Grenoble) 56 (2006), 2093–2113.
- [9] B. Adamczewski, Y. Bugeaud et F. Luca, *Sur la complexité des nombres algébriques*, C. R. Math. Acad. Sci. Paris 339 (2004), 11–14.
- [10] B. Adamczewski and J. Cassaigne, *Diophantine properties of real numbers generated by finite automata*, Compos. Math. 142 (2006), 1351–1372.
- [11] J.-P. Allouche, J. L. Davison, M. Queffélec, and L. Q. Zamboni, *Transcendence of Sturmian or morphic continued fractions*, J. Number Theory 91 (2001), 39–66.
- [12] J.-P. Allouche and J. Shallit, *Automatic Sequences: Theory, Applications, Generalizations*. Cambridge University Press, Cambridge, 2003.
- [13] A. Baker, *Continued fractions of transcendental numbers*, Mathematika 9 (1962), 1–8.
- [14] A. Baker, *On Mahler’s classification of transcendental numbers*, Acta Math. 111 (1964), 97–120.
- [15] V. Berthé, C. Holton and L. Q. Zamboni, *Initial powers of Sturmian words*, Acta Arith. 122 (2006), 315–347.
- [16] Y. Bugeaud, *Approximation by algebraic numbers*. Cambridge Tracts in Mathematics 160, Cambridge, 2004.
- [17] J. L. Davison, *Quasi-Periodic Continued Fractions*, J. Number Theory 127 (2007), 272–282.
- [18] F. Durand, *Linearly recurrent subshifts have a finite number of non-periodic subshift factors*, Ergod. Th. & Dynam. Sys. 20 (2000), 1061–1078.
- [19] J.-H. Evertse, *An improvement of the quantitative Subspace theorem*, Compositio Math. 101 (1996), 225–311.
- [20] N. I. Feldman and Yu. V. Nesterenko, *Number Theory. IV. Transcendental Numbers*. Encyclopaedia of Mathematical Sciences 44. Springer–Verlag, Berlin, 1998.
- [21] S. Ferenczi and Ch. Mauduit, *Transcendence of numbers with a low complexity expansion*, J. Number Theory 67 (1997), 146–161.
- [22] S. Fischler, *Palindromic Prefixes and Episturmian Words*, J. Combin. Theory Ser. A 113 (2006), 1281–1304.
- [23] K. Mahler, *Zur Approximation der Exponentialfunktionen und des Logarithmus. I, II*, J. reine angew. Math. 166 (1932), 118–150.
- [24] E. Maillet, *Introduction à la théorie des nombres transcendants et des propriétés arithmétiques des fonctions*. Gauthier-Villars, Paris, 1906.
- [25] O. Perron, *Die Lehre von den Kettenbrüchen*. Teubner, Leipzig, 1929.

- [26] M. Queffélec, *Transcendance des fractions continues de Thue–Morse*, J. Number Theory 73 (1998), 201–211.
- [27] D. Ridout, *Rational approximations to algebraic numbers*, Mathematika 4 (1957), 125–131.
- [28] K. F. Roth, *Rational approximations to algebraic numbers*, Mathematika 2 (1955), 1–20; corrigendum, 168.
- [29] W. M. Schmidt, *Norm form equations*, Ann. of Math. 96 (1972), 526–551.
- [30] W. M. Schmidt, *Diophantine Approximation*. Lecture Notes in Mathematics 785, Springer, 1980.
- [31] W. M. Schmidt, *The subspace theorem in Diophantine approximation*, Compositio Math. 69 (1989), 121–173.
- [32] Th. Schneider, *Einführung in die transzendenten Zahlen*. Springer–Verlag, Berlin–Göttingen–Heidelberg, 1957.

Boris Adamczewski
 CNRS, Université de Lyon, Université Lyon 1
 Institut Camille Jordan
 Bât. Braconnier, 21 avenue Claude Bernard
 69622 VILLEURBANNE Cedex (FRANCE)
 Boris.Adamczewski@math.univ-lyon1.fr

Yann Bugeaud
 Université Louis Pasteur
 U. F. R. de mathématiques
 7, rue René Descartes
 67084 STRASBOURG Cedex (FRANCE)
 bugeaud@math.u-strasbg.fr