

HYPERTRANSCENDENCE AND LINEAR DIFFERENCE EQUATIONS

BORIS ADAMCZEWSKI, THOMAS DREYFUS, AND CHARLOTTE HARDOUIN

ABSTRACT. After Hölder proved his classical theorem about the Gamma function, there has been a whole bunch of results showing that solutions to linear difference equations tend to be hypertranscendental (*i.e.*, they cannot be solution to an algebraic differential equation). In this paper, we obtain the first complete results for solutions to general linear difference equations associated with the shift operator $x \mapsto x + h$ ($h \in \mathbb{C}^*$), the q -difference operator $x \mapsto qx$ ($q \in \mathbb{C}^*$ not a root of unity), and the Mahler operator $x \mapsto x^p$ ($p \geq 2$ integer). The only restriction is that we constrain our solutions to be expressed as (possibly ramified) Laurent series in the variable x with complex coefficients (or in the variable $1/x$ in some special case associated with the shift operator). Our proof is based on the parametrized difference Galois theory initiated by Hardouin and Singer. We also deduce from our main result a general statement about algebraic independence of values of Mahler functions and their derivatives at algebraic points.

1. INTRODUCTION

Number theorists have to face the following challenge. On the one hand, the fields of rational and algebraic numbers are too poor, while, on the other hand, the fields of real and complex numbers seem far too large (uncountable). An attempt to classify transcendental numbers leads to the introduction of the ring of periods \mathcal{P} (see [KZ01]) and to the following classification:

$$\mathbb{Q} \subset \overline{\mathbb{Q}} \subset \mathcal{P} \subset \mathbb{C}.$$

The main features of \mathcal{P} are that most of classical mathematical constants (*e.g.*, π , $\log 2$, $\zeta(3)$, $\Gamma(1/3)^3 \dots$)* belong to it, and that \mathcal{P} is a countable set whose elements contain only a finite amount of information and can be identified in an algorithmic way. After the pioneering works of Cauchy and Riemann, analytic and holomorphic functions figure prominently in mathematics. As already observed by Hilbert [Hil02], their study gives rise to a similar difficulty: the fields of rational and algebraic functions are too limited, but the ring $\mathbb{C}[[x]]$ is huge. In order to overcome this deficiency, he suggested to classify transcendental functions according to whether they are *holonomic*

Date: July 10, 2020.

2010 *Mathematics Subject Classification.* 39A06, 12H05.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program under the Grant Agreement No 648132, and from the ANR De rerum natura project (ANR-19-CE40-0018).

*However, there should be some exceptions. For instance, it is conjectured that e is not a period.

(*i.e.*, satisfy a linear differential equation with polynomial coefficients), *differentially algebraic* (*i.e.*, satisfy an algebraic differential equation), or not. Hence, we obtain the following classification:

$$(1.1) \quad \text{Rat} \subset \text{Alg} \subset \text{Hol} \subset \text{Dif} \subset \mathbb{C}[[x]].$$

Functions that are not differentially algebraic are said to be *hypertranscendental*. Note that, if we restrict our attention to power series with algebraic coefficients, the rings Hol and Dif become countable, and, again, their elements contain only a finite amount of information and can be identified in an algorithmic way. These rings play, in the setting of holomorphic functions, a role similar to the one played by \mathcal{P} in the setting of complex numbers. The classification (1.1) is also significant in enumerative combinatorics. To some extent, the nature of a generating series reflects the underlying structure of the objects it counts (see the discussion in [BM06]). An appealing example, where all cases of (1.1) appear at once, is given by the study of generating series associated with lattice walks (see, for instance, [DHRS18] and the references therein).

Hilbert also observed that the class of differentially algebraic functions misses important functions coming from number theory. Here are few examples. The very first comes from a classical result of Hölder [Höl87] stating that the gamma function $\Gamma(x)$ is hypertranscendental. Furthermore, it follows from the identity

$$\zeta(x) = 2(2\pi)^{x-1}\Gamma(1-x)\sin\left(\frac{\pi x}{2}\right)\zeta(1-x)$$

that the Riemann zeta function is also hypertranscendental. In an other direction, Moore [Moo96] has shown that $f_1(x) = \sum_{n \geq 0} x^{2^n}$ is hypertranscendental. A result reproved and extended later by Mahler [Mah30a] in connection with the so-called Mahler's method in transcendental number theory. More recently, Hardouin and Singer [HS08] proved hypertranscendence for some q -hypergeometric series, such as

$$f_2(x) := \sum_{n=0}^{\infty} \frac{(1-a)^2(1-aq)^2 \cdots (1-aq^{n-1})^2}{(1-q)^2(1-q^2)^2 \cdots (1-q^n)^2} x^n,$$

where $q \in \mathbb{C}^*$ is not a root of unity, $a \notin q^{\mathbb{Z}}$ and $a^2 \in q^{\mathbb{Z}}$. Most proofs about hypertranscendence turn out to be based on the fact that the function under consideration satisfies a functional equation of a different type (*i.e.*, not differential). For instance, the three results we just mentioned are respectively based on the following linear difference equations:

$$\Gamma(x+1) = x\Gamma(x), \quad f_1(x^2) = f_1(x) - x,$$

and

$$f_2(q^2x) - \frac{2ax-2}{a^2x-1}f_2(qx) + \frac{x-1}{a^2x-1}f_2(x) = 0.$$

Broadly speaking, satisfying both a linear difference equation and an algebraic differential equation would enforce too much symmetry for a transcendental function. In this paper, we follow this motto:

(A) *A solution to a linear difference equation should be either rational or hypertranscendental.*

Of course, this must be taken with a pinch of salt. For instance, $\log(x)$, $\exp(x)$, and the Jacobi theta function $\theta_q(x) = \sum_{n \in \mathbb{Z}} q^{-n(n-1)/2} x^n$ are all differentially algebraic, but they also satisfy the following simple linear difference equations:

$$\log(x^p) = p \log x, \quad \exp(x+1) = e \exp(x), \quad \theta_q(qx) = qx\theta_q(x).$$

Nevertheless, we obtain in this paper the first complete results about hypertranscendence of solutions to general linear difference equations associated with the shift operator $x \mapsto x+h$ ($h \in \mathbb{C}^*$), the q -difference operator $x \mapsto qx$ ($q \in \mathbb{C}^*$ not a root of unity), and the Mahler operator $x \mapsto x^p$ ($p \geq 2$ integer). The only restriction is that we constrain our solutions to be expressed as (possibly ramified) Laurent series in the variable x (or in the variable $1/x$ in some special case associated with the shift operator). For a precise statement, we refer the reader to our main result, Theorem 1.2, which confirms guess (A) for these three operators.

1.1. Statement of our main result. All along this paper, our framework consists in a tower of field extensions $\mathbb{C} \subset K \subset F_0 \subset F$ with the following properties.

- The field K is equipped with an automorphism ρ and a derivation ∂^\dagger .
- The automorphism ρ extends to F_0 and F .
- The derivation ∂ extends to F , but F_0 is not necessarily closed under ∂ .

Our aim is to prove that, if $f \in F_0$ is solution to a linear ρ -equation with coefficients in K , then either $f \in K$ or f is hypertranscendental over K . Specifically, we consider the following four situations, which we refer to as Cases **S**₀, **S**_∞, **Q**, and **M**, respectively.

Case S₀. In this case, we let $K = \mathbb{C}(x)$, $\partial = \frac{d}{dx}$, and ρ denote the automorphism of K defined by

$$\rho(f(x)) = f(x+h), \quad h \in \mathbb{C}^*.$$

The automorphism ρ and the derivation ∂ naturally extend to the field $F := \text{Mer}(\mathbb{C})$ of meromorphic functions on \mathbb{C} . Let $K_0 := \mathbb{C}(x, \exp(*x))$ denote the field generated over $\mathbb{C}(x)$ by the functions $\exp(\ell x)$, $\ell \in \mathbb{C}$. In this setting, we let F_0 be any field extension of $\mathbb{C}(x)$ on which ρ extends and satisfying the following three conditions.

- (i) $F_0 \subset \text{Mer}(\mathbb{C})$.
- (ii) $\{f \in F_0 \mid \rho(f) = f\} = \mathbb{C}$.
- (iii) $F_0 \cap K_0 = \mathbb{C}(x)$.

Remark 1.1. For $h = 1$, let us see that we can choose $F_0 = \mathbb{C}(x, \Gamma(x))$, where $\Gamma(x)$ is the Gamma function. The functional equation $\Gamma(x+1) = x\Gamma(x)$ shows that ρ extends to F_0 and it is well-known that $\Gamma \in \text{Mer}(\mathbb{C})$. Thus, (i) holds. Now, let $\beta \in F_0$ be such that $\rho(\beta) = \beta$. Writing β as a rational function in Γ with coefficients in $\mathbb{C}(x)$, and then using the functional equation satisfied by Γ and the fact that Γ is transcendental over $\mathbb{C}(x)$,

[†]A derivation ∂ on K is a map from K into itself satisfying $\partial(a+b) = \partial(a) + \partial(b)$ and $\partial(ab) = \partial(a)b + a\partial(b)$ for every $a, b \in K$.

we can deduce that $\beta \in \mathbb{C}$. Hence (ii) holds. Finally, to prove (iii) it is sufficient to show that Γ is transcendental over K_0 . The latter property follows from classical asymptotics showing that, for every $\alpha(x) \in K_0$, we have $\lim_{x \rightarrow +\infty} |\Gamma(x)/\alpha(x)| = +\infty$. Thus, Case \mathbf{S}_0 of Theorem 1.2 is a generalization of Hölder's theorem.

Case \mathbf{S}_∞ . In this case, we consider K , ρ , and ∂ as in Case \mathbf{S}_0 . We note that the automorphism ρ and the derivation ∂ naturally extend to the field of Laurent series $\mathbb{C}((x^{-1}))$, and we set $F_0 = F = \mathbb{C}((x^{-1}))$.

Case \mathbf{Q} . In this case, we let $K = \bigcup_{j \geq 1} \mathbb{C}(x^{1/j})$ denote the field of ramified rational functions. We also use the notation $\mathbb{C}(x^{1/*})$ for this field. Given $q \in \mathbb{C}^*$ that is not a root of unity, we let ρ denote the automorphism of K defined by

$$\rho(f(x)) = f(qx), \quad \partial = x \frac{d}{dx},$$

and we let $F_0 = F = \bigcup_{j \geq 1} \mathbb{C}((x^{1/j}))$ denote the field of Puiseux series. We also use the notation $\mathbb{C}((x^{1/*}))$ for this field.

Case \mathbf{M} . In this case, we let $K = \mathbb{C}(x^{1/*})$ and, given a natural number $p \geq 2$, we let ρ denote the automorphism of K defined by

$$\rho(f(x)) = f(x^p),$$

and we set $\partial = x \frac{d}{dx}$ and $F_0 = F = \mathbb{C}((x^{1/*}))$.

In all cases, ∂ is a derivation on K and F . An element f in F is said to be *differentially algebraic* or ∂ -algebraic over K if there exists $n \in \mathbb{N}$ such that the functions $f, \partial(f), \dots, \partial^{(n)}(f)$ are algebraically dependent over K . Otherwise, f is said to be *hypertranscendental* over K . Our main result reads as follow.

Theorem 1.2. *Let K , F_0 , and ρ be defined as in Cases \mathbf{S}_0 , \mathbf{S}_∞ , \mathbf{Q} , and \mathbf{M} , and let n be a positive integer. Let $f \in F_0$ be solution to the ρ -linear difference equation of order n*

$$(1.2) \quad \rho^n(y) + a_{n-1}\rho^{n-1}(y) + \dots + a_1\rho(y) + a_0y = 0,$$

where $a_0, \dots, a_{n-1} \in K$. Then either $f \in K$ or f is hypertranscendental over K .

On the road to Theorem 1.2, there is already a whole bunch of partial results and a number of them are used in our proof. These results can roughly be divided into three different types.

- Those concerned with affine equations of order one, that is with equations of the form $\rho(y) = ay + b$. References include [Höl87, Moo96, Mah30a, Nis84, Ran92, Ish98, Har08, HS08, Ngu12].
- Those concerned with equations whose difference Galois group is large (*e.g.*, simple, semisimple, reductive). References include [HS08, DHR18, DHR16, AS17, ADR18]. In particular, such results have nice applications to equations of order 2.

- Those dealing with general equations, but reaching only nonholonomicity instead of hypertranscendence (see Theorem 1.3 and Remark 1.4 below). References include [Ram92, BG93, Béz94, SS19].

Among these results, we quote the following one, which may be considered as a first step towards a proof of Theorem 1.2 (see Remark 1.4 for precise attribution).

Theorem 1.3. *In each of the cases \mathbf{S}_0 , \mathbf{S}_∞ , \mathbf{Q} , and \mathbf{M} , the following property holds. Any element of F_0 that satisfies both a ρ -linear equation and a ∂ -linear equation with coefficients in K belongs to K .*

Remark 1.4. Theorem 1.3 can be rephrased by saying that, if f is as in Theorem 1.2, then either $f \in K$ or f is not holonomic. Note that recently the authors of [SS19] give a uniform proof for all cases of Theorem 1.3, see Corollaries 3 and 5 therein. In each case, we also mention the original reference. Case \mathbf{S}_0 is due to Bézivin and Gramain [BG93]. Case \mathbf{S}_∞ is due to Schäfke and Singer [SS19, Corollary 5]. Furthermore, in Cases \mathbf{Q} and \mathbf{M} , a change of variable of the form $z = x^{1/\ell}$, $\ell \in \mathbb{N}^*$, can be used to reduce the situation to the case where $K = \mathbb{C}(x)$ and $F_0 = \mathbb{C}((x))$. In the latter situation, Case \mathbf{Q} is due to Ramis [Ram92], while Case \mathbf{M} is due to Bézivin [Béz94].

1.2. Strategy of proof. Elements of the rings Rat , Alg , and Hol are well-understood. We have recurrence formulas for their coefficients, precise asymptotics, and a good understanding of their finitely many singularities. In contrast, our knowledge about differentially algebraic functions is much more limited (see, for instance, the survey [Rub89]). In fact, such functions can behave quite wildly. For example, the function $\sum x^{n^2}$ is differentially algebraic and admits the unit circle as a natural boundary[‡]. Thus, proving hypertranscendence required more involved tools, and is substantially more difficult than proving irrationality, transcendence, or nonholonomicity. In this direction, Hardouin and Singer [HS08] built a new Galois theory of difference equations, which is designed to measure the differential dependencies among solutions to linear difference equations. In particular, it applies to hypertranscendence. We refer the reader to [DV12] and [Har16] for an introduction to this topic. Our proof of Theorem 1.2 relies on this theory.

Let us briefly describe our strategy. Since f satisfies a linear difference equation, it is classical to associate with this equation its difference Galois group, which is a linear algebraic group that encodes the algebraic relations between the solutions to this equation. The more involved parametrized Galois theory developed in [HS08] attaches to any linear difference equation, a geometric object, *the parametrized Galois group*, whose structure encodes the algebraic relations satisfied by the solutions and their derivatives. This is not a linear algebraic group anymore, but a linear differential group (see Section 2.2). Nevertheless, both groups are strongly related and, viewed on a suitable field extension, the latter is Zariski dense in the former (see Proposition 2.5). In this framework, we can conclude that f is hypertranscendental

[‡]Incidentally, this example shows that the "outrageous conjecture" suggested at the end of [Rub89] is false. Indeed, it follows from a theorem of Nesterenko that $\sum 2^{-n^2}$ is transcendental.

when the corresponding parametrized Galois group is *large*. Furthermore, there are several results, culminating in [AS17], showing that if the classical difference Galois group is large, then both groups are the same when viewed on a suitable field extension. We emphasize that such results follow a strategy initiated in [HS08] and developed further in [DHR18]. It combines a fundamental result about classification of differential algebraic subgroups of semisimple algebraic groups by Cassidy [Cas72, Cas89], parametrized Galois correspondence, and Theorem 1.3.

We prove Theorem 1.2 by induction on the order n of the ρ -linear equation satisfied by f . We first prove the result for affine equations of order one. Then we show that, iterating the associated linear system if necessary, one can reduce the situation to the case where the associated difference Galois group G is connected. Now, if G acts irreducibly on \mathbb{C}^n , we show how to reduce the situation to the case where G is semisimple. In that case, the difference Galois group is large enough and a recent result of Arreche and Singer [AS17] allows us to conclude that f is hypertranscendental. Finally, if G acts reducibly on \mathbb{C}^n , we assume that f is differentially algebraic and we show how to construct from f a differentially algebraic function g that satisfies a linear equation of smaller order. Furthermore, this construction ensures that $g \in K$ if and only if $f \in K$. Applying the induction assumption to g , we deduce that $f \in K$, as wanted. Thus, once the case of order one equation is solved, the proof really makes a dichotomy between *irreducible* and *reducible* difference Galois groups.

1.3. An application to number theory. At the end of the 1920s, Mahler [Mah29, Mah30b, Mah30a] introduced a new method for proving transcendence and algebraic independence of values at algebraic points of analytic functions satisfying different types of functional equations associated with the operator $x \mapsto x^p$. Mahler's method aims at transferring results about algebraic independence over $\mathbb{Q}(x)$ of such functions to the algebraic independence over $\overline{\mathbb{Q}}$ of their values at algebraic points. A major result in this direction is Nishioka's theorem [Nis90], which provides an analogue of the Siegel-Shidlovskii theorem for linear Mahler systems. Let us also mention that significant progress in this theory has been made recently [Phi15, AF17, AF18a, AF18b]. Combining Case **M** of Theorem 1.2 with Nishioka's theorem and following the line of argument given in the proof of Proposition 4.1 in [AF18b], we can deduce the following general result. We recall that $f(x) \in \overline{\mathbb{Q}}[[x]]$ is a *p-Mahler function* if there exist polynomials $a_0(x), \dots, a_n(x) \in \overline{\mathbb{Q}}[x]$, not all zero, such that

$$a_0(x)f(x) + a_1(x)f(x^p) + \dots + a_n(x)f(x^{p^n}) = 0.$$

We simply say that f is a Mahler function if it is p -Mahler for some integer $p \geq 2$. According to Theorem 1.2, a Mahler function is thus either rational or hypertranscendental. We recall that an irrational Mahler function is meromorphic on the open unit disc and admits the unit circle as a natural boundary.

Theorem 1.5. *Let $f(x) \in \overline{\mathbb{Q}}[[z]]$ be a Mahler function that is not rational, let r be a positive integer, and let \mathcal{K} be a compact subset of the open unit*

disc. Then, for all but finitely many algebraic numbers $\alpha \in \mathcal{K}$, the complex numbers $f(\alpha), f'(\alpha), \dots, f^{(r)}(\alpha)$ are algebraically independent over $\overline{\mathbb{Q}}$.

Results of this type have been first obtained by Mahler [Mah30a] for solutions to affine order one equations. The main feature of Theorem 1.5 is that it applies to *any* irrational Mahler function. It is also almost best possible in the sense that one cannot avoid the possibility of finding infinitely many exceptions in the whole open unit disc. For instance, the transcendental 2-Mahler function

$$f(z) = \prod_{n=0}^{\infty} (1 - 3z^{2^n})$$

vanishes at all algebraic numbers α such that $\alpha^{2^n} = 1/3$ for some integer $n \geq 0$. This shows that, even for $r = 1$, the exceptional set can depend on \mathcal{K} . However, if \mathcal{K} is fixed, we do not know whether the exceptional set always remains finite when r tends to infinity.

1.4. Organization of the paper. This article is organized as follows. In Section 2, we provide a short introduction to difference Galois theory and to parametrized difference Galois theory, following [vdPS97, HS08]. In Section 3, we describe, for each Cases \mathbf{S}_0 , \mathbf{S}_∞ , \mathbf{Q} , and \mathbf{M} , the suitable framework that is needed to use the parametrized difference Galois theory of [HS08]. Several auxiliary results are gathered in Section 4. They concern algebraic groups, the behavior of the difference Galois group when considered over different field extensions, and Picard-Vessiot extensions associated with iterated systems. Finally, Section 5 is devoted to the proof of Theorem 1.2.

Acknowledgments. We warmly thank Michael F. Singer for his careful reading of an earlier version of this paper and for making many useful suggestions. We are also grateful to Daniel Bertrand for his comments.

2. DIFFERENCE AND PARAMETRIZED DIFFERENCE GALOIS THEORIES

In this section, we provide a short introduction to difference Galois theory and to parametrized difference Galois theory.

2.1. Difference Galois theory. We first recall some notation, as well as classical results, concerning difference Galois theory. We refer the reader to [vdPS97] for more details.

2.1.1. Notation in operator algebra. A difference ring is a pair (R, ρ) where R is a ring and ρ is a ring automorphism of R . If R is a field, then (R, ρ) is called a difference field or a ρ -field. An ideal I of R such that $\rho(I) \subset I$ is called a difference ideal or a ρ -ideal. We say that the difference ring (R, ρ) is simple if the only ρ -ideals of R are $\{0\}$ and R . Two difference rings (R_1, ρ_1) and (R_2, ρ_2) are isomorphic if there exists a ring isomorphism φ between R_1 and R_2 such that $\varphi \circ \rho_1 = \rho_2 \circ \varphi$. A difference ring (S, ρ') is a difference ring extension of (R, ρ) if S is a ring extension of R and if $\rho'|_R = \rho$. In this case, we usually keep on denoting ρ' by ρ . When R is a ρ -field, we say that S is a R - ρ -algebra. Two difference ring extensions (R_1, ρ) and (R_2, ρ) of the

difference ring (R, ρ) are isomorphic over (R, ρ) if there exists a difference ring isomorphism φ from (R_1, ρ) to (R_2, ρ) such that $\varphi|_R = \text{Id}_R$. The ring of constants of the difference ring (R, ρ) is defined by

$$R^\rho := \{f \in R \mid \rho(f) = f\}.$$

If R^ρ is a field, it is called the field of constants. If there is no risk of confusion, we usually simply say that R , instead of (R, ρ) , is a difference ring (or a difference field, or a difference ring extension...).

2.1.2. *Difference equations and linear difference systems.* Let (K, ρ) be a difference field. A linear ρ -equation of order n over K is an equation of the form

$$(2.1) \quad \mathcal{L}(y) := \rho^n(y) + a_{n-1}\rho^{n-1}(y) + \cdots + a_0y = 0,$$

with $a_0, \dots, a_{n-1} \in K$. If $a_0 \neq 0$, this relation can be written in matrix form as

$$(2.2) \quad \rho Y = A_{\mathcal{L}} Y$$

where

$$A_{\mathcal{L}} := \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \ddots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 \\ 0 & 0 & \cdots & 0 & 1 \\ -a_0 & -a_1 & \cdots & \cdots & -a_{n-1} \end{pmatrix} \in \text{GL}_n(K).$$

The matrix $A_{\mathcal{L}}$ is called the companion matrix associated with Equation (2.1). It is often more convenient to use the notion of linear difference system, that is of system of the form

$$(2.3) \quad \rho(Y) = AY, \text{ with } A \in \text{GL}_n(K).$$

We recall that two difference systems $\rho(Y) = AY$ and $\rho(Y) = BY$ with $A, B \in \text{GL}_n(K)$ are said to be *equivalent over K* if there exists a gauge transformation $T \in \text{GL}_n(K)$ such that $B = \rho(T)AT^{-1}$. In that case, $\rho(Y) = AY$ if and only if $\rho(TY) = B(TY)$.

Remark 2.1. By [HS99, Theorem B.2.], if K contains a nonperiodic element with respect to ρ , then the cyclic vector lemma holds, and any linear difference system is equivalent to one given by a companion matrix associated with some linear equation. Let L be a ρ -field extension of K and let $(f_0, f_1, \dots, f_{n-1})^\top \in L^n$ be a nonzero solution to a linear system $\rho(Y) = AY$, with $A \in \text{GL}_n(K)$. Then each coordinate f_j satisfies some linear ρ -equation over K of order at most n .

2.1.3. *Picard-Vessiot rings and difference Galois groups.* A *Picard-Vessiot ring* for (2.3) over a difference field (K, ρ) of characteristic zero is a K - ρ -algebra satisfying the following three properties.

- (1) There exists $U \in \text{GL}_n(R)$ such that $\rho(U) = AU$. Such a matrix U is called a *fundamental matrix*.
- (2) R is generated as a ring over K by the coordinates of U and by $\det(U)^{-1}$, that is $R = K[U, \det(U)^{-1}]$.
- (3) R is a simple difference ring.

A Picard-Vessiot ring is not always an integral domain. However, it is a direct sum of integral domains which are transitively permuted by ρ . More precisely, by [vdPS97, Corollary 1.16], there exist $r \geq 1$ and orthogonal idempotent elements e_0, \dots, e_{r-1} , such that

$$R = e_0R \oplus \dots \oplus e_{r-1}R,$$

where e_iR is an integral domain and $\rho(e_i) = e_{i+1 \bmod r}$. We recall that Picard-Vessiot rings always exist. Indeed, it is sufficient to endow the polynomial K -algebra $K[X, \frac{1}{\det(X)}]$ with a structure of K - ρ -algebra by setting $\rho(X) = AX$. Then, for any maximal ρ -ideal \mathfrak{M} of $K[X, \frac{1}{\det(X)}]$, the quotient $R = K[X, \frac{1}{\det(X)}]/\mathfrak{M}$ is a Picard-Vessiot ring. However, this construction does not guarantee that the ring of ρ -constant has not grown. This justifies the introduction of the more convenient notion of Picard-Vessiot extension.

A *Picard-Vessiot extension* \mathcal{Q} for (2.3) over a difference field (K, ρ) of characteristic zero is a K - ρ -algebra \mathcal{Q} satisfying the following properties.

- (1) There exists $U \in \mathrm{GL}_n(\mathcal{Q})$ such that $\rho(U) = AU$. Such a matrix U is called a fundamental matrix.
- (2) \mathcal{Q} is a *pseudofield* extension of K , that is there exist a positive integer r , orthogonal idempotent elements e_0, \dots, e_{r-1} of \mathcal{Q} , and a field extension L of K such that $\mathcal{Q} = e_0L \oplus \dots \oplus e_{r-1}L$ and $\rho(e_i) = e_{i+1 \bmod r}$.
- (3) \mathcal{Q} is the smallest pseudofield extension of K containing U .
- (4) $\mathcal{Q}^\rho = K^\rho$.

When \mathcal{Q} is a field, we say that \mathcal{Q} is a *Picard-Vessiot field extension*. By [vdPS97, §1.1], if K^ρ is algebraically closed, then there exists a Picard-Vessiot extension and two Picard-Vessiot extensions are isomorphic as K - ρ -algebras. The relation between Picard-Vessiot-rings and Picard-Vessiot-extensions is given by the following proposition, which is a straightforward consequence of [OW15, Proposition 2.5 and Corollary 2.6].

Proposition 2.2. *If K^ρ is an algebraically closed field of characteristic zero, then the following properties hold.*

- Let \mathcal{Q} be a Picard-Vessiot extension over K and let us define $R := K[U, \frac{1}{\det(U)}] \subset \mathcal{Q}$, where U is a fundamental matrix. Then R is a Picard-Vessiot ring.
- If R is a Picard-Vessiot ring over K then the total quotient ring[§] of R is a Picard-Vessiot extension.

From now on, we assume that K is a ρ -field such that $\mathbf{k} := K^\rho$ is an algebraically closed field of characteristic zero. Let \mathcal{Q} be a Picard-Vessiot extension over K . The *difference Galois group* $\mathrm{Gal}(\mathcal{Q}/K)$ of (2.3) over K is defined as the group of K - ρ -algebra automorphisms of \mathcal{Q} :

$$\mathrm{Gal}(\mathcal{Q}/K) := \{\sigma \in \mathrm{Aut}(\mathcal{Q}/K) \mid \rho \circ \sigma = \sigma \circ \rho\}.$$

For any fundamental matrix $U \in \mathrm{GL}_n(\mathcal{Q})$, an easy computation shows that $U^{-1}\sigma(U) \in \mathrm{GL}_n(\mathbf{k})$ for all $\sigma \in \mathrm{Gal}(\mathcal{Q}/K)$. By [vdPS97, Theorem 1.13],

[§]We recall that, given a ring R , its total quotient ring is defined as the localization of R at the multiplicative set formed by all nonzero divisors of R .

the faithful representation

$$\begin{aligned} \mathrm{Gal}(\mathcal{Q}/K) &\rightarrow \mathrm{GL}_n(\mathbf{k}) \\ \sigma &\mapsto U^{-1}\sigma(U) \end{aligned}$$

identifies $\mathrm{Gal}(\mathcal{Q}/K)$ with a linear algebraic subgroup $G \subset \mathrm{GL}_n(\mathbf{k})$. Choosing another fundamental matrix of solutions U leads to a conjugate representation.

2.1.4. Torsor and algebraic relations. A fundamental result in difference Galois theory ([vdPS97, Theorem 1.13]) says that the Picard-Vessiot ring R is the coordinate ring of a $\mathrm{Gal}(\mathcal{Q}/K)$ -torsor over K . Thereby, the difference Galois group controls the algebraic relations satisfied by the solutions to the underlying linear system. As a corollary of this structure of $\mathrm{Gal}(\mathcal{Q}/K)$ -torsor, one obtains the fundamental equality

$$\dim_{\mathbf{k}}\mathrm{Gal}(\mathcal{Q}/K) = \mathrm{trdeg}_K \mathcal{Q},$$

where we let $\mathrm{trdeg}_K \mathcal{Q}$ denote the transcendence degree of the field extension L/K . Here, the field L is defined as in (2) of the definition of a Picard-Vessiot extension.

2.1.5. The Galois correspondence. The Galois correspondence for linear difference systems can be summarized as follows (see [vdPS97, Theorem 1.29]).

Theorem 2.3. *Let \mathcal{Q} be a Picard-Vessiot extension over K . Let \mathcal{R} denote the set of K - ρ -algebras F such that $F \subset \mathcal{Q}$ and such that every nonzero divisor of F is a unit of F . Let \mathcal{G} denote the set of algebraic subgroups of $\mathrm{Gal}(\mathcal{Q}/K)$. Then the following properties hold.*

- For any $F \in \mathcal{R}$, the set

$$G(\mathcal{Q}/F) := \{\sigma \in \mathrm{Gal}(\mathcal{Q}/K) \mid \sigma(f) = f, \quad \forall f \in F\}$$

is an algebraic subgroup of $\mathrm{Gal}(\mathcal{Q}/K)$.

- For any $H \in \mathcal{G}$, the set

$$\mathcal{Q}^H := \{f \in \mathcal{Q} \mid \sigma(f) = f, \quad \forall \sigma \in H\}$$

belongs to \mathcal{R} .

- The maps

$$\begin{array}{ccc} r : \mathcal{R} \rightarrow \mathcal{G} & & g : \mathcal{G} \rightarrow \mathcal{R} \\ F \mapsto G(\mathcal{Q}/F) & \text{and} & H \mapsto \mathcal{Q}^H \end{array}$$

are each other's inverses.

Remark 2.4. In the case where R is an integral domain, \mathcal{Q} is a difference field, and Theorem 2.3 provides a correspondence between the difference subfields of \mathcal{Q} containing K , on the one hand, and the algebraic subgroups of $\mathrm{Gal}(\mathcal{Q}/K)$, on the other hand.

2.2. Parametrized difference Galois theory. We recall here some notation and results concerning parametrized difference Galois theory. We refer the reader to [HS08] for more details.

2.2.1. *Notation in operator algebra.* A differential ring is a pair (R, δ) where R is a ring and δ is a derivation on R , that is a homomorphism of the additive group from R into itself satisfying the Leibniz rule:

$$\delta(ab) = \delta(a)b + a\delta(b).$$

If R is a field, then (R, δ) is called a differential field or a δ -field. The ring of constants of R is defined by

$$R^\delta = \{f \in R \mid \delta(f) = 0\}.$$

Let L be a δ -ring extension of a δ -ring R . Given a subset S of L , we let $R\{S\}_\delta$ denote the smallest R - δ -algebra generated by S . A field K endowed with a structure of both difference and differential field is called a (ρ, δ) -field if ρ and δ commute, that is if

$$\rho \circ \delta(f) = \delta \circ \rho(f)$$

for every $f \in K$.

The study of algebraic structures of difference and differential fields is the main object of the so-called differential and difference algebra. We refer the interested reader to the founding books of Kolchin ([Kol73]) for further details concerning differential algebra and to the book of Cohn for further details concerning difference algebra ([Coh65]). It is worth mentioning that these two setting are very different. In order to avoid many analogous definitions, we use the following convention: we add an operator prefix to an algebraic attribute to signify the compatibility of the algebraic structure with respect to the operator. For instance, a (ρ, δ) -ring is a ring equipped with ρ and δ , a (ρ, δ) -field extension $K \subset L$ is a field extension such that the fields L and K are (ρ, δ) -fields and such that the action of ρ and δ are compatible with the field extension, and so on.

2.2.2. *(ρ, δ) -Picard-Vessiot ring and the parametrized difference Galois group.* Let \tilde{K} be a (ρ, δ) -field and let $A \in \text{GL}_n(\tilde{K})$. We consider the difference system

$$(2.4) \quad \rho(Y) = AY.$$

A (ρ, δ) -Picard-Vessiot ring for (2.4) over \tilde{K} is a (ρ, δ) -ring extension \tilde{R} of \tilde{K} satisfying the following three properties.

- (1) There exists $U \in \text{GL}_n(\tilde{R})$ such that $\rho(U) = AU$. Such a matrix U is called a fundamental matrix.
- (2) \tilde{R} is generated as a δ -ring over \tilde{K} by the coordinates of U and by $\det(U)^{-1}$, that is $\tilde{R} = \tilde{K}\{U, \det(U)^{-1}\}_\delta$.
- (3) \tilde{R} is a simple (ρ, δ) -ring, that is the only (ρ, δ) -ideals of \tilde{R} are $\{0\}$ and \tilde{R} .

A (ρ, δ) -Picard-Vessiot ring is not always an integral domain but it is a direct sum of integral domains closed under δ and transitively permuted by ρ . A construction similar to that of Section 2.1.3 shows that (ρ, δ) -Picard-Vessiot rings always exist. Again, this construction does not ensure that the ring of ρ -constants has not grown. This justifies the introduction of the more convenient notion of (ρ, δ) -Picard-Vessiot extension.

A (ρ, δ) -Picard-Vessiot extension $\tilde{\mathcal{Q}}$ for (2.4) over \tilde{K} is a \tilde{K} - ρ - δ -algebra satisfying the following properties.

- (1) There exists $U \in \mathrm{GL}_n(\tilde{\mathcal{Q}})$ such that $\rho(U) = AU$. Such a matrix U is called a fundamental matrix.
- (2) $\tilde{\mathcal{Q}}$ is a pseudo δ -field extension of \tilde{K} , that is there exist a positive integer r , orthogonal idempotent elements e_0, \dots, e_{r-1} of $\tilde{\mathcal{Q}}$, and \tilde{L} a δ -field extension of \tilde{K} such that $\tilde{\mathcal{Q}} = e_0\tilde{L} \oplus \dots \oplus e_{r-1}\tilde{L}$ and $\rho(e_i) = e_{i+1 \bmod r}$.
- (3) $\tilde{\mathcal{Q}}$ is equal to $\tilde{K}\langle U \rangle$, that is the smallest pseudo δ -field extension of \tilde{K} containing U .
- (4) $\tilde{\mathcal{Q}}^\rho = \tilde{K}^\rho$.

When $\tilde{\mathcal{Q}}$ is a field, we say that $\tilde{\mathcal{Q}}$ is a (ρ, δ) -Picard-Vessiot field extension. By [Wib12, Corollary 9], if \tilde{K}^ρ is algebraically closed, then there exists a (ρ, δ) -Picard-Vessiot extension $\tilde{\mathcal{Q}}$ for (2.4) over \tilde{K} . The smallest \tilde{K} - (ρ, δ) -subalgebra $\tilde{K}\{U, \frac{1}{\det(U)}\}_\delta$ of $\tilde{\mathcal{Q}}$ generated by a fundamental matrix and the inverse of its determinant is a (ρ, δ) -Picard-Vessiot ring. However, in order to ensure the uniqueness of (ρ, δ) -Picard-Vessiot rings up to \tilde{K} - (ρ, δ) -algebra isomorphisms, one needs the field \tilde{K}^ρ to be δ -closed ([HS08, Proposition 6.16]). We recall that a differential field (L, δ) is called *differentially closed* or δ -closed if, for every set of δ -polynomials \mathcal{F} , the system of δ -equations $\mathcal{F} = 0$ has a solution in some δ -field extension of L if and only if it has a solution in L . Note that a δ -closed field is algebraically closed. Differentially closed fields are huge and, for instance, none of the function fields K introduced in Section 1.1 satisfies the assumption that K^ρ is differentially closed. Nonetheless, working with δ -closed fields allows us to simplify many arguments. Thus, we will embed K into a (ρ, δ) -field \tilde{K} whose field of ρ -constants is δ -closed, and we will use some descent argument to go back to K if necessary (see Sections 3 and 4.2).

From now on, we let \tilde{K} denote a (ρ, δ) -field whose field of ρ -constants $\tilde{C} := \tilde{K}^\rho$ is δ -closed. Let $\tilde{\mathcal{Q}}$ be a (ρ, δ) -Picard-Vessiot extension over \tilde{K} . The *parametrized difference Galois group* $\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$ of (2.4) over $(\tilde{K}, \rho, \delta)$, also called the (ρ, δ) -Galois group, is defined as the group of \tilde{K} - (ρ, δ) -algebra automorphisms of $\tilde{\mathcal{Q}}$:

$$\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K}) := \left\{ \sigma \in \mathrm{Aut}(\tilde{\mathcal{Q}}/\tilde{K}) \mid \rho \circ \sigma = \sigma \circ \rho \text{ and } \delta \circ \sigma = \sigma \circ \delta \right\}.$$

The parametrized difference Galois group $\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$ is a geometric object that encodes the differential algebraic relations between the solutions to (2.4). See [HS08, Proposition 6.24] for more details. Roughly speaking, the larger $\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$, the fewer δ -algebraic relations over \tilde{K} that hold among the elements of $\tilde{\mathcal{Q}}$.

For any fundamental matrix $U \in \mathrm{GL}_n(\tilde{\mathcal{Q}})$, an easy computation shows that $U^{-1}\sigma(U) \in \mathrm{GL}_n(\tilde{C})$ for any element $\sigma \in \mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$. By [HS08, Proposition 6.18], the faithful representation

$$\begin{aligned} \mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/K) &\rightarrow \mathrm{GL}_n(\tilde{C}) \\ \sigma &\mapsto U^{-1}\sigma(U) \end{aligned}$$

identifies $\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$ with a *linear differential algebraic subgroup* $H \subset \mathrm{GL}_n(\tilde{C})$, i.e. a subgroup of $\mathrm{GL}_n(\tilde{C})$ that can be defined as the vanishing set of polynomial δ -equations with coefficients in \tilde{C} . Furthermore, choosing another fundamental matrix U leads to a conjugate representation.

2.2.3. Comparison between difference Galois groups and parametrized difference Galois groups. Let \tilde{R} denote a (ρ, δ) -Picard-Vessiot ring for the system (2.4) over \tilde{K} , let $\tilde{\mathcal{Q}}$ denote the associated (ρ, δ) -Picard-Vessiot extension, and let $U \in \mathrm{GL}_n(\tilde{R})$ be a fundamental matrix of solutions. By [HS08, Proposition 6.21], $R := \tilde{K}[U, \det(U)^{-1}]$ is a Picard-Vessiot ring for (2.4) over \tilde{K} . Let \mathcal{Q} denote the Picard-Vessiot extension corresponding to R . Then we have $\mathcal{Q} \subset \tilde{\mathcal{Q}}$ and one can identify $\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$ with a subgroup of $\mathrm{Gal}(\mathcal{Q}/\tilde{K})$ by restricting the elements of $\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$ to \mathcal{Q} . The following result provides a fundamental link between classical difference Galois theory and its parametrized counterpart, see [HS08, Proposition 6.21].

Proposition 2.5. *The parametrized difference Galois group $\mathrm{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K})$ is a Zariski-dense subgroup of $\mathrm{Gal}(\mathcal{Q}/\tilde{K})$.*

3. THE PARAMETRIZED FRAMEWORK

Let us go back to our $K, F_0, F, \rho, \partial$ corresponding to Cases $\mathbf{S}_0, \mathbf{S}_\infty, \mathbf{Q}$, and \mathbf{M} (see Section 1.1). We first observe that, in each case, $K^\rho = \mathbb{C}$ is algebraically closed and K has characteristic zero. Hence the theory of [vdPS97] described in Section 2.1 applies. In this section, we describe some suitable field extensions \tilde{K} of K , and \tilde{C} of $C = F^\rho$, that allow us to apply the parametrized difference Galois theory of [HS08] described in Section 2.2.

We first recall that any differential field K has a differential closure, *i.e.* a differentially closed field extension that can be embedded in every differentially closed field extension of K . In Case \mathbf{S}_0 , we set $\delta := \partial$, and we let \tilde{C} denote a δ -closure of C_h , the subfield of $\mathrm{Mer}(\mathbb{C})$ formed by those meromorphic functions that are h -periodic. Note that C_h is also a δ -field for ρ and δ commute. Sometimes, we will use the notation \tilde{C}_h instead of \tilde{C} . In Cases \mathbf{S}_∞ and \mathbf{Q} , we also set $\delta := \partial$, and we let $\tilde{\mathbb{C}}$ denote a δ -closure of \mathbb{C} and we set $\tilde{C} := \tilde{\mathbb{C}}$. By [DHR18, Lemma 2.3], in these three cases, the field

$$\tilde{K} := \mathrm{Frac}(K \otimes_{\mathbb{C}} \tilde{C})$$

is a (ρ, δ) -field extension of K such that $\tilde{K}^\rho = \tilde{C}$.

Case \mathbf{M} is a bit more tricky. Indeed, in order to apply the theory of [HS08], we need to consider a derivation δ on \tilde{K} such that δ and ρ commute.

Following [DHR18], we let \log denote a transcendental element over K , and we define $\delta = \log \times \partial$ to be the derivation that acts on $K(\log)$ by

$$\delta(\log) = \log \quad \text{and} \quad \delta(x^a) = ax^a \log .$$

Then we define a structure of ρ -field on $K(\log)$ by setting $\rho(\log) = p \log$. We observe that $K(\log)^\rho = \mathbb{C}$ and that ρ and δ commute. Finally, we consider $\widetilde{\mathbb{C}}$ a δ -closure of \mathbb{C} such that ρ acts trivially on $\widetilde{\mathbb{C}}$. We set $\widetilde{C} := \widetilde{\mathbb{C}}$ and by [DHR18, Lemma 2.3], the field

$$\widetilde{K} := \text{Frac}(K(\log) \otimes_{\mathbb{C}} \widetilde{C}) ,$$

is a (ρ, δ) -field such that $\widetilde{K}^\rho = \widetilde{C}$.

In all cases, K is a ρ -field of characteristic zero such that $K^\rho = \mathbb{C}$, F is a ρ -field extension of K with $C := F^\rho$, and \widetilde{K} is a (ρ, δ) -field of characteristic zero, such that $\widetilde{K}^\rho = \widetilde{C}$, where \widetilde{C} is differentially closed. Furthermore, ρ and δ commute. The following table summaries the different frameworks we consider in this paper.

Case	F	K	C	∂	δ	\widetilde{C}	\widetilde{K}
\mathbf{S}_0	$\mathcal{M}er(\mathbb{C})$	$\mathbb{C}(x)$	C_h	$\frac{d}{dx}$	$\frac{d}{dx}$	\widetilde{C}_h	$\widetilde{C}_h(x)$
\mathbf{S}_∞	$\mathbb{C}((x^{-1}))$	$\mathbb{C}(x)$	\mathbb{C}	$\frac{d}{dx}$	$\frac{d}{dx}$	$\widetilde{\mathbb{C}}$	$\widetilde{\mathbb{C}}(x)$
\mathbf{Q}	$\mathbb{C}((x^{1/*}))$	$\mathbb{C}(x^{1/*})$	\mathbb{C}	$x \frac{d}{dx}$	$x \frac{d}{dx}$	$\widetilde{\mathbb{C}}$	$\widetilde{\mathbb{C}}(x^{1/*})$
\mathbf{M}	$\mathbb{C}((x^{1/*}))$	$\mathbb{C}(x^{1/*})$	\mathbb{C}	$x \frac{d}{dx}$	$\log \times x \frac{d}{dx}$	$\widetilde{\mathbb{C}}$	$\widetilde{\mathbb{C}}(x^{1/*})(\log)$

Remark 3.1. Note that the field of δ -constants of C is the algebraically closed field \mathbb{C} . Then, by [CS07, Lemma 9.3], the field of δ -constants of \widetilde{C} is also \mathbb{C} . At some places, we will consider some iteration of the operator ρ . In Case \mathbf{S}_0 , it might happen that we start with the ρ -field extension $\widetilde{K} = \widetilde{C}_h(x)$ of K , and then consider this extension as a ρ^r -field extension. Note that $\widetilde{C}_h \neq \widetilde{C}_{rh}$. Nonetheless, \widetilde{C}_h is a ρ^r -constant field and all results of Section 2 can be applied to the (ρ^r, δ) -field \widetilde{K} .

4. AUXILIARY RESULTS

In this section, we let (K, ρ) be a difference field of characteristic zero and we consider a ρ -linear difference system

$$(4.1) \quad \rho(Y) = AY, \quad \text{with } A \in \text{GL}_n(K),$$

where $\mathbf{k} := K^\rho$ is an algebraically closed field.

4.1. Irreducible and reducible difference Galois groups. The strategy for proving Theorem 1.2 is very different according to whether the difference Galois group associated with the difference equation over K satisfied by f is irreducible or not. By irreducible, we mean that the group acts irreducibly on \mathbf{k}^n , see Definition 4.1. We point out that this is different from saying that an algebraic group, viewed as an algebraic variety, is irreducible.

Definition 4.1. Let $G \subset \text{GL}_n(\mathbf{k})$ be a group. We say that G is *irreducible* if it acts irreducibly on \mathbf{k}^n , that is, if the only \mathbf{k} -vector subspaces of \mathbf{k}^n invariant by G are $\{0\}$ and \mathbf{k}^n . We say that G is *reducible* otherwise. When

G is an irreducible group, we say that G is *imprimitive* if there exist an integer $r \geq 2$, and V_1, \dots, V_r , some \mathbf{k} -vector spaces satisfying the following conditions.

- (i) $\mathbf{k}^n = V_1 \oplus \dots \oplus V_r$.
- (ii) For every $g \in G$, the mapping $V_i \mapsto g(V_i)$ is a permutation of the set $\{V_1, \dots, V_r\}$.

We say that G is *primitive* otherwise.

The next result shows that a difference Galois group that is irreducible and imprimitive cannot be connected. We recall that an algebraic group G is connected if it has no proper open subgroup with respect to the Zariski topology. The *identity component* of G is the connected component of G containing the identity.

Lemma 4.2. *Let $G \subset \mathrm{GL}_n(\mathbf{k})$ be an algebraic group. If G is connected and irreducible, then G is primitive.*

Proof. Let us assume by contradiction that G is imprimitive. Then there exist an integer $r \geq 2$ and some nonzero \mathbb{C} -vector-spaces $V_1, \dots, V_r \subsetneq \mathbf{k}^n$ such that $\mathbf{k}^n = V_1 \oplus \dots \oplus V_r$. Furthermore, the action induced by every element of G on \mathbf{k}^n is a permutation of the sets $\{V_1, \dots, V_r\}$. Let G_{V_1} denote the stabilizer of V_1 in G . This is an algebraic subgroup of G . We claim that it has finite index in G . Let us assume by contradiction that there are an infinite number of cosets $g_i G_{V_1}$ of G_{V_1} in G . Since the group of permutations of a set with r elements is finite, there exist two distinct cosets $g_1 G_{V_1}$ and $g_2 G_{V_1}$ such that g_1 and g_2 induce the same permutation of the V_i 's. In that case, $g_2^{-1} g_1$ stabilizes V_1 and thus belongs to G_{V_1} , providing a contradiction. This proves the claim. Now, since G_{V_1} is a closed subgroup of G of finite index, it contains the identity component of G . The latter is equal to G for G is assumed to be connected. This means that V_1 is invariant under the action of G . Hence, G is reducible. This provides a contradiction. \square

Now, let us recall the definition of a difference module, see [vdPS97, §1.4].

Definition 4.3. A K -difference module \mathcal{M} is a pair (M, Σ) where M is a finite-dimensional K -vector space and Σ is an additive endomorphism of M such that $\Sigma(\lambda Y) = \rho(\lambda)\Sigma(Y)$ for every $Y \in M$ and $\lambda \in K$.

Let $A \in \mathrm{GL}_n(K)$. The K -difference module \mathcal{M}_A attached to the system $\rho(Y) = AY$ is $\mathcal{M}_A = (K^n, \Sigma_A)$, where $\Sigma_A(Y) = A^{-1}\rho(Y)$. Given a monic linear difference operator $\mathcal{L} = a_0 y + a_1 \rho(y) + \dots + \rho^n(y)$ with coefficients a_i in K and $a_0 \neq 0$, one can consider the corresponding linear difference system $\rho(Y) = A_{\mathcal{L}} Y$, where $A_{\mathcal{L}}$ is the companion matrix of \mathcal{L} . If K contains a nonperiodic element with respect to ρ , then [HS99, Theorem B.2] asserts that any K -difference module \mathcal{M} is isomorphic to $\mathcal{M}_{A_{\mathcal{L}}}$ for some monic linear difference operator \mathcal{L} . This is equivalent to the existence of a so-called cyclic vector in \mathcal{M} , that is, a nonzero element $e \in M$ such that the vectors $e, \Sigma(e), \dots, \Sigma^{n-1}(e)$ form a K -basis of M .

The next lemma characterizes linear difference systems whose difference Galois group is reducible.

Lemma 4.4. *Let $G \subset \mathrm{GL}_n(\mathbf{k})$ be the difference Galois group of (4.1) over K and let r be an integer with $0 < r < n$. The following statements are equivalent.*

- *There exists a G -submodule of \mathbf{k}^n [¶] of dimension r over \mathbf{k} .*
- *There exists a difference submodule of dimension r in the K -difference module \mathcal{M}_A associated with (4.1).*
- *There exists $T \in \mathrm{GL}_n(K)$ such that*

$$\rho(T)AT^{-1} = \begin{pmatrix} B_1 & B_2 \\ 0 & B_3 \end{pmatrix}$$

with $B_1 \in \mathrm{GL}_r(K)$.

In particular, G is reducible if and only if the statements above hold for some r , with $0 < r < n$. Furthermore, if the matrix A in (4.1) is the companion matrix of a difference operator \mathcal{L} , then \mathcal{L} admits a nontrivial right factor if and only if G is reducible.

Proof. Since the category of K -difference modules is a Tannakian category, the first two statements are nothing else than the usual Tannakian equivalence, see [And01, Théorème 3.2.1.1]. Completing a K -basis of a difference submodule \mathcal{M} in \mathcal{M}_A of dimension r , $0 < r < n$, to a K -basis $\underline{e} = (e_1, \dots, e_n)$ of K^n , shows that the matrix of Σ in the basis \underline{e} is of the form $B = \begin{pmatrix} B_1 & B_2 \\ 0 & B_3 \end{pmatrix}$, with $B_1 \in \mathrm{GL}_r(K)$. Let $T \in \mathrm{GL}_n(K)$ denote the matrix associated with the change of basis from the canonical basis $\underline{f} = (f_1, \dots, f_n)$ to \underline{e} . Then

$$BT\underline{f} = B\underline{e} = \Sigma(\underline{e}) = \Sigma(T\underline{f}) = \rho(T)A\underline{f}.$$

This proves the equivalence between the second and the third statements.

Now, let us assume that A is the companion matrix of some difference operator \mathcal{L} and that there exists a nontrivial difference submodule \mathcal{N} of \mathcal{M}_A . Let e be a cyclic vector of \mathcal{M}_A such that \mathcal{L} is the minimal monic linear difference operator annihilating e . Since \mathcal{N} is a nontrivial difference submodule of \mathcal{M}_A , then $\dim_K \mathcal{M}_A / \mathcal{N} < n$ and there exists a nontrivial monic linear difference operator \mathcal{L}_1 of order smaller than or equal to $\dim_K \mathcal{M}_A / \mathcal{N}$ such that $\mathcal{L}_1 e \in \mathcal{N}$. The element $\mathcal{L}_1 e$ is nonzero by minimality of \mathcal{L} . Since \mathcal{N} is a difference submodule of \mathcal{M}_A there exists a monic linear difference operator \mathcal{L}_2 of order smaller than or equal to $\dim_K \mathcal{N}$ such that $\mathcal{L}_2 \mathcal{L}_1(e) = 0$. By minimality of \mathcal{L} , this proves that $\mathcal{L} = \mathcal{L}_2 \mathcal{L}_1$, where \mathcal{L}_1 has order smaller than n . Conversely, if there is a nontrivial factorization $\mathcal{L} = \mathcal{L}_2 \mathcal{L}_1$, then the difference module corresponding to \mathcal{L}_1 is a nontrivial difference submodule of \mathcal{M} . Hence G is reducible. \square

4.2. Going up and down. In this section, we let K, F_0, ρ be defined as in Section 1.1, and we let \tilde{K} be defined as in Section 3. The two following lemmas compare the difference Galois groups over K and over \tilde{K} .

We recall that a connected linear algebraic group G is said to be *semisimple* if it is of positive dimension and its only solvable (or, equivalently, Abelian) connected closed normal subgroup is the trivial group.

[¶] \mathbf{k}^n is a G -module via $G \subset \mathrm{GL}_n(\mathbf{k})$.

Lemma 4.5. *Let $A \in \mathrm{GL}_n(K)$ and let G (resp. \tilde{G}) be the Galois group of $\rho(Y) = AY$ over K (resp. \tilde{K}). The following properties hold.*

- In Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{Q} , we have $\tilde{G} = G(\tilde{C})$.
- In Case \mathbf{M} , if G is semisimple, then $\tilde{G} = G(\tilde{C})$.

Proof. In Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{Q} , the field extension \tilde{K} of K is obtained by extension of the field of constants \mathbb{C} to \tilde{C} . Then, [CHS08, Corollary 2.5] gives that the difference Galois group over \tilde{K} is just the extension of G to the \tilde{C} -points.

Let us deal with Case \mathbf{M} . Let \mathcal{G} denote the difference Galois group over $K(\log)$. As previously, we infer from [CHS08, Corollary 2.5] that $\tilde{G} = \mathcal{G}(\tilde{C})$. Thus, it remains to prove that $\mathcal{G} = G$ when G is semisimple. By [DHR18, Proposition 1.6], we can see \mathcal{G} as a subgroup of G such that one of the following two situations occurs.

- (1) $\mathcal{G} = G$.
- (2) \mathcal{G} is a normal subgroup of G and $G/\mathcal{G} \simeq \mathbf{G}_m$ the multiplicative group (\mathbb{C}^*, \cdot) .

By [Mil17, Corollary 21.50], a semisimple algebraic group is perfect, that is equal to its derived subgroup, see [Mil17, Definition 12.45]. Thus, a semisimple algebraic group has no nontrivial abelian quotient. Hence, (2) cannot occur. We deduce that $G = \mathcal{G}$, which ends the proof. \square

Lemma 4.6. *Let $A \in \mathrm{GL}_n(K)$ and let G (resp. \tilde{G}) be the Galois group of $\rho(Y) = AY$ over K (resp. \tilde{K}). The group G is irreducible if and only if the group \tilde{G} is irreducible.*

Proof. In Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{Q} , Lemma 4.5 ensures that $\tilde{G} = G(\tilde{C})$ and the result follows directly.

Let us deal with Case \mathbf{M} . We infer from [CHS08, Corollary 2.5] that $\tilde{G} = \mathcal{G}(\tilde{C})$, where we let \mathcal{G} denote the Galois group of the system over $K(\log)$. Thus, it remains to prove that \mathcal{G} is irreducible if and only if G is irreducible. As seen in the proof of Lemma 4.5, \mathcal{G} is a subgroup of G . Thus, \mathcal{G} is reducible as soon as G is reducible. Conversely, let us assume that \mathcal{G} is reducible. By the cyclic vector lemma over K , the linear difference system (4.1) is equivalent to an equation

$$\mathcal{L} := \rho^n + a_{n-1}\rho^{n-1} + \cdots + a_0 = 0,$$

with $a_i \in K$ and $a_0 \neq 0$. Since \mathcal{G} is reducible, Lemma 4.4 gives that \mathcal{L} admits a factorization over $K(\log)$. Multiplying \mathcal{L} by some nonzero element if necessary, we can get rid of the denominators. Then, there exists $\alpha \in K[\log]$ such that

$$\alpha \mathcal{L} = (b_k \rho^k + b_{k-1} \rho^{k-1} + \cdots + b_0)(c_{n-k} \rho^{n-k} + c_{n-k-1} \rho^{n-k-1} + \cdots + c_0),$$

where $b_i, c_i \in K[\log]$, $b_k b_0 c_{n-k} c_0 \neq 0$, and $0 < k < n$. Rewriting the equation in terms of powers of \log , one finds

$$\sum_{i \geq i_0} \log^i(\alpha_i \mathcal{L}) = \left(\sum_{\kappa \geq \kappa_0} \log^\kappa \mathcal{L}_\kappa \right) \left(\sum_{j \geq j_0} \log^j \mathcal{D}_j \right),$$

where $\mathcal{L}_\kappa, \mathcal{D}_j$ are linear difference operators over K , $\alpha_i \in K$, $\alpha_{i_0} \in K^*$, and $\mathcal{L}_{\kappa_0} := \sum_{r=0}^t \gamma_r \rho^r$ and \mathcal{D}_{j_0} are nonzero. Using the fact that \log is transcendental over K and $\rho \log = p \log \rho$, we deduce that $i_0 = \kappa_0 + j_0$. Considering the terms in \log^{i_0} in both sides of the factorization, we deduce that $\mathcal{L} = \frac{1}{\alpha_{i_0}} \mathcal{M} \mathcal{D}_{j_0}$ where $\mathcal{M} = \sum_{r=0}^t \gamma_r p^{j_0 r} \rho^r$, leading to a nontrivial factorization of \mathcal{L} over K . By Lemma 4.4, we obtain that G is reducible, as wanted. \square

Now, we prove the following descent lemma.

Lemma 4.7. *For F_0, K and \tilde{K} defined as in Section 3, we have*

$$F_0 \cap \tilde{K} = K.$$

Proof. Let us first consider Case **Q**. Let $a \in F_0 \cap \tilde{C}(x^{1/*})$. There exists a positive integer r such that $a \in \mathbb{C}((x^{1/r})) \cap \tilde{C}(x^{1/r})$. It follows that there exists a positive integer N such that $x^N a \in \mathbb{C}[[x^{1/r}]]$. Thus, $x^N a$ is a formal power series in $x^{1/r}$ with complex coefficients that represents a rational fraction with coefficients in \tilde{C} . This property can be characterized by the vanishing of its associated Kronecker-Hankel determinants (see, for instance, [Sal63, p. 5]). This condition does not depend on the field of coefficients, so we deduce that $x^N a \in \mathbb{C}(x^{1/r})$. Hence, $a \in K$.

The proof of Case **S_∞** is entirely similar. For Case **M**, if $a \in \mathbb{C}((x^{1/*})) \cap \tilde{C}(x^{1/*})(\log)$, we use the fact that the logarithm is transcendental over $\mathbb{C}((x^{1/*}))$ to conclude that $a \in \mathbb{C}((x^{1/*})) \cap \tilde{C}(x^{1/*})$. Arguing as above, we can show that $a \in K$.

Now, let us deal with Case **S₀**. Let $a \in F_0 \cap \tilde{C}(x)$. Since $F_0 \subset \text{Mer}(\mathbb{C})$, one can find $x_0 \in \mathbb{C}$ such that $a \in \mathbb{C}[[x - x_0]]$. Once again, using Kronecker-Hankel determinants, one can deduce that $a \in K$. \square

4.3. Iterating the difference operator. For every positive integer ℓ , let us set

$$A_{[\ell]} := \rho^{\ell-1}(A) \times \cdots \times A \in \text{GL}_n(K).$$

Note that if U is a fundamental matrix of $\rho(Y) = AY$, then it is also a fundamental matrix of $\rho^\ell(Y) = A_{[\ell]}Y$. In other words, a vector solution to the system $\rho(Y) = AY$ is also a solution to the iterated system $\rho^\ell(Y) = A_{[\ell]}Y$. In this section, we show that, replacing the original system by a suitable iteration, we can reduce the situation to the more convenient case where the corresponding Galois group is connected and the Picard-Vessiot extension is a field.

This iteration process might introduce some new constants. The following lemma shows that this cannot happen when the field of ρ -constants is algebraically closed.

Lemma 4.8. *Let L be a ρ -field such that $L^\rho = \mathbf{k}$, and r be a positive integer. Then, any ρ^r -constant of L is algebraic of degree less than or equal to r over \mathbf{k} . In particular, if \mathbf{k} is algebraically closed then $L^{\rho^r} = \mathbf{k}$.*

Proof. Let $a \in L$ be a ρ^r -constant. Then the polynomial $P(X) = (X - a) \cdots (X - \rho^{r-1}(a))$ belongs to $L^\rho[X] = \mathbf{k}[X]$. This proves that a is algebraic over \mathbf{k} of degree less than or equal to r . \square

One of the main difficulty of difference algebra is the control of the algebraic difference field extensions. This is the source of many technical difficulties and connected to the fact that the theory ACFA is unstable (see [CH99]). The difference fields (K, ρ) introduced in Section 1.1 have the following very strong property with respect to finite difference field extensions, which will allow us to bypass these difficulties.

Lemma 4.9. *Let r be a positive integer. In each of the cases \mathcal{S}_0 , \mathcal{S}_∞ , \mathcal{Q} , and \mathcal{M} , there are no proper ρ^r -field extension of K of finite degree.*

Proof. The results in [Hen97, Hen98, Roq18] cover all cases. \square

We are now ready to prove the following result.

Proposition 4.10. *Let (K, ρ) be defined as in Section 1.1 and let L be a ρ -field extension of K such that $L^\rho = \mathbb{C}$. Let $(f_1, \dots, f_n)^\top \in L^n$ be a nonzero solution to (4.1). Let G be the difference Galois group of (4.1) over K . There exists a positive integer r such that the following properties hold.*

- (a) *There exists a Picard-Vessiot field extension \mathcal{Q} for $\rho^r(Y) = A_{[r]}Y$ over (K, ρ^r) , with a fundamental solution matrix U having $(f_1, \dots, f_n)^\top$ as first column.*
- (b) *The difference Galois group $G_r := \text{Gal}(\mathcal{Q}/K)$ of $\rho^r(Y) = A_{[r]}Y$ over (K, ρ^r) coincides with the identity component of G and is therefore connected.*

Though this result can be easily deduced from the proof of [DHR16, Proposition 4.6] in the particular case where the parametric operator is the identity, we find it more convenient for the reader to include the proof below.

Proof. (a) Since $K(f_1, \dots, f_n)^\rho \subset L^\rho = \mathbb{C}$, there exists a Picard-Vessiot ring R_0 for $\rho(Y) = AY$ over $K(f_1, \dots, f_n)$. Since $f_1, \dots, f_n \in R_0$, there exists a fundamental matrix U for R_0 having $(f_1, \dots, f_n)^\top$ as first column. Let \mathcal{Q}_0 be the total quotient ring of R_0 . By Proposition 2.2, we find $\mathcal{Q}_0^\rho = \mathbb{C}$.

Set $R := K[U, 1/\det(U)]$. The difference ring (R, ρ) is a subring of the Picard-Vessiot ring (R_0, ρ) . Since any element of R that is a zero divisor in R_0 is a zero divisor in R , see for instance [Roq18, Lemma 5], the total quotient ring \mathcal{Q} of R embeds in \mathcal{Q}_0 and consequently is a pseudofield (see [Wib10, Lemma 1.3.4]). Consequently $\mathbb{C} \subset \mathcal{Q}^\rho \subset \mathcal{Q}_0^\rho = \mathbb{C}$. Then $\mathcal{Q}^\rho = \mathbb{C}$, and (\mathcal{Q}, ρ) is a Picard-Vessiot extension for $\rho(Y) = AY$ over (K, ρ) . Since $K^\rho = \mathbb{C}$ is algebraically closed, the uniqueness of Picard-Vessiot extensions above K allows to conclude that $\text{Gal}(\mathcal{Q}/K) = G$. We have proved that we can embed any solution in L in a Picard-Vessiot extension above K .

By Proposition 2.2, R is a Picard-Vessiot ring for $\rho(Y) = AY$ over (K, ρ) . Let e_0, \dots, e_{r-1} denote the orthogonal idempotent relative to its ring structure. Note that $K[f_1, \dots, f_n]$ is an integral domain that is closed under ρ . For $i = 0, \dots, r-1$, let $\phi_i : R \rightarrow e_i R$ denote the projection from R to $e_i R$. Note that $\phi_i(x) = x e_i$. Let us prove that the restriction to $K[f_1, \dots, f_n]$ of this projection is injective for all $i \in \{0, \dots, r-1\}$. Suppose to the contrary that there exist a nonzero element $x \in K[f_1, \dots, f_n]$ and $i \in \{0, \dots, r-1\}$ such that $\phi_i(x) = 0$. Since ρ permutes the integral domains $e_i R$, we find that $x \rho(x) \cdots \rho^{r-1}(x) = 0$. The latter equality provides a contradiction with the fact that $K[f_1, \dots, f_n]$ is an integral domain which is closed under the injective morphism ρ . Thus, $e_i K[f_1, \dots, f_n] \subset e_i R$ is a copy of $K[f_1, \dots, f_n]$ in

$e_i R$. Since $K^\rho = \mathbb{C}$ is algebraically closed, Lemma 4.8 implies that $K^{\rho^r} = \mathbb{C}$. By [vdPS97, Lemma 1.26], $(e_i R, \rho^r)$ is a Picard-Vessiot ring for the ρ^r -system $\rho^r(Y) = A_{[r]}Y$ and $e_i R$ is an integral domain. Its total quotient ring \mathcal{Q}_1 is a field and, since K^{ρ^r} is algebraically closed, \mathcal{Q}_1 is a Picard-Vessiot field extension by Proposition 2.2. By construction, \mathcal{Q}_1 contains f_1, \dots, f_n , and we can thus choose a fundamental matrix having $(f_1, \dots, f_n)^\top$ as first column.

(b) By [Roq18, Theorem 12], $G_r = \text{Gal}(\mathcal{Q}_1/K)$ is a normal algebraic subgroup of G and the quotient G/G_r is finite. To conclude, it remains to prove that G_r is connected. Let G_r° denote its identity component. The Galois correspondence, see Theorem 2.3, gives that the ρ^r -field $\mathcal{Q}_1^{G_r^\circ}$ is a finite extension of K . Lemma 4.9 implies that $\mathcal{Q}_1^{G_r^\circ} = K$ and, applying the Galois correspondence again, we deduce that $G_r^\circ = G_r$. \square

Remark 4.11. The proof of (b) shows that, if G is connected, then, for every positive integer ℓ , the Galois group of $\rho^\ell(Y) = A_{[\ell]}Y$ over K coincides with G .

In the parametrized framework, we have the following similar result.

Proposition 4.12. *Let L be a (ρ, δ) -field extension of \tilde{K} with $L^\rho = \tilde{C}$, and let $(f_1, \dots, f_n)^\top \in L^n$ be a nonzero solution to (4.1). Then there exist a positive integer r and a parametrized Picard-Vessiot field extension $\tilde{\mathcal{Q}}$ for $\rho^r(Y) = A_{[r]}Y$ over \tilde{K} , with a fundamental matrix U having $(f_1, \dots, f_n)^\top$ as first column.*

Proof. The proof is similar to Proposition 4.10 and [DHR18, Lemma 3.7], just noting that, by Lemma 4.8, $\tilde{K}^{\rho^r} = \tilde{C}$ for \tilde{C} is algebraically closed. \square

The following result is obtained by combining the two previous propositions.

Corollary 4.13. *Let $(f_1, \dots, f_n)^\top \in F_0^n$ be a nonzero solution to (4.1) and let G be the difference Galois group of (4.1). Then, there exists a positive integer r such that*

- (a) *There exists a (ρ^r, δ) -Picard-Vessiot field extension $\tilde{\mathcal{Q}}$ for $\rho^r(Y) = A_{[r]}Y$ with a fundamental matrix U having $(f_1, \dots, f_n)^\top$ as first column over \tilde{K} (where $\tilde{K} = \overline{C_{sh}(x)}$ for some positive integer s dividing r in Case \mathbf{S}_0).*
- (b) *The difference Galois group of $\rho^r(Y) = A_{[r]}Y$ over K coincides with the identity component of G .*

Proof. By Proposition 4.10, there exists a positive integer s such that the difference Galois group of $\rho^s(Y) = A_{[s]}Y$ is equal to the identity component of G . In order to apply Proposition 4.12 to the system $\rho^s(Y) = A_{[s]}Y$, we need to embed F_0 into a (ρ^s, δ) -field extension L of \tilde{K} with $L^{\rho^s} = \tilde{C}$. We proceed as follows.

- Case \mathbf{S}_0 . We let F_1 denote the smallest (ρ^s, δ) -subfield of $\text{Mer}(\mathbb{C})$ containing F_0 and C_{sh} . Since $\text{Mer}(\mathbb{C})^{\rho^s} = C_{sh}$, we find that $F_1^{\rho^s} = C_{sh}$. Let \tilde{C} denote a δ -closure of C_{sh} , considered as a constant ρ^s -field. The field extensions \tilde{C} and F_1 are linearly disjoint above C_{sh}

(see [Wib10, Lemma 1.1.6]). Thereby, their compositum L is the fraction field of $\tilde{C} \otimes_{C_{sh}} F_1$ (see [Bou90, A.V.13]). Arguing as in the proof of [DHR18, Lemma 2.3], we obtain that L has the required properties.

- Cases \mathbf{S}_∞ and \mathbf{Q} . We let L be the fraction field of $F_0 \otimes_C \tilde{C}$.
- Case \mathbf{M} . We let L be the field $\tilde{C}((x^{1/*}))(\log)$.

Applying Proposition 4.12 to the system $\rho^s(Y) = A_{[s]}Y$ over \tilde{K} , one can perform a second iteration to obtain a system $\rho^{sl}(Y) = A_{[sl]}Y$ satisfying (a). By Remark 4.11, the difference Galois group of $\rho^{sl}(Y) = A_{[sl]}Y$ over K is the identity component of G . This ends the proof. \square

5. PROOF OF THEOREM 1.2

This section is devoted to the proof of our main result. Before proving Theorem 1.2 in full generality, we first consider the following two particular cases.

- The function f is solution to an inhomogeneous equation of order one.
- The difference Galois group of the equation associated with f is both connected and irreducible.

All along this section, we keep on with the notation of Sections 1.1 and 3.

5.1. Affine order one equations. Various hypertranscendence criteria for solutions to inhomogeneous order one equations have already been obtained, see for instance [Ran92, Ish98, HS08, Ngu12]. We first deduce from these criteria the following result.

Proposition 5.1. *If $f \in F_0$ is solution to an equation of the form $\rho(f) = af + b$, with $a, b \in K$, then either f is ∂ -transcendental over K or f belongs to K .*

Proof of Proposition 5.1. Let us first note that if $a = 0$, then $f = \rho^{-1}(b) \in K$. We can thus assume that $a \neq 0$. Furthermore, we observe that, in Cases \mathbf{Q} and \mathbf{M} , we can use a change of variable of the form $z = x^{1/\ell}$ to reduce the situation to the case where $K = \mathbb{C}(x)$. From now on, we thus assume that $K = \mathbb{C}(x)$. Let us also assume that f is ∂ -algebraic over K . It remains to prove that $f \in K$.

We first note that Case \mathbf{M} corresponds to [Ran92, Théorème 5.2]. Now, let us consider Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{Q} , which are partially covered by [HS08]. We recall the following definition from [HS08, §6.1] (see also [vdPS97, §2]). We say that $\alpha \in K$ is *standard* if, for all positive integers ℓ , α and $\rho^\ell(\alpha)$ have no common zero or pole. By [HS08, Lemma 6.2], there exists a standard element a^* and a nonzero $e \in K$ such that $a = a^* \rho(e)/e$. It follows that f/e is solution to the equation

$$\rho(f/e) = a^* f/e + b/\rho(e).$$

Set $g = f/e$ and $\tilde{b} = b/\rho(e)$. Since $e \in K$, it suffices to prove that $g \in K$. Let us note that $g = f/e$ is ∂ -algebraic over K .

The same argument as in [ADR18, Lemma 3.6] shows that there exists a nonzero δ -algebraic solution of $\rho(y) = a^*y + b/\rho(e)$ in a (ρ, δ) -Picard-Vessiot

field extension of \tilde{K} . In Case \mathbf{S}_0 , [HS08, Proposition 3.9] implies that $a^* \in \mathbb{C}$ and there exists $h \in K$ such that $\tilde{b} = \rho(h) - a^*h$. In Case \mathbf{S}_∞ , the same argument as in the proof of [HS08, Proposition 3.9] also yields the same conclusion. We obtain that

$$\begin{aligned}\rho(g) &= a^*g + \tilde{b} \\ &= a^*g + \rho(h) - a^*h.\end{aligned}$$

Hence $\rho(g - h) = a^*(g - h)$. If $g = h$, then $g \in K$ and we are done. Thus, we can assume that $g \neq h$. In that case, setting $\tilde{f} = g - h$, we obtain $\rho(\partial(\tilde{f})) = a^*\partial(\tilde{f})$. It follows that $\rho(\partial(\tilde{f})/\tilde{f}) = \partial(\tilde{f})/\tilde{f}$, and thus $\partial(\tilde{f})/\tilde{f} \in \mathbb{C} \subset K$. Since $\tilde{f} \in F_0$ satisfies both a linear ρ -equation and a linear ∂ -equation with coefficients in K , Theorem 1.3 implies that $\tilde{f} \in K$. Since $h \in K$, we deduce that $g \in K$, as expected.

In Case \mathbf{Q} , the same argument as in the proof of [HS08, Proposition 3.10] shows that $a^* = cx^\alpha$ for some $c \in \mathbb{C}^*$ and $\alpha \in \mathbb{Z}$, and one of the following conditions holds.

- (a) $a^* = q^r$ for some $r \in \mathbb{Z}$, and $\tilde{b} = \rho(h) - a^*h + dx^r$ for some $h \in K$ and $d \in \mathbb{C}$.
- (b) $a^* \notin q^{\mathbb{Z}}$ and $\tilde{b} = \rho(h) - a^*h$ for some $h \in K$.

In Case (a), we deduce that

$$(5.1) \quad \rho\left(\frac{g-h}{x^r}\right) = \frac{g-h}{x^r} + dq^{-r}.$$

Since $(g-h)/x^r \in F_0$, we can consider its expansion in Puiseux series. Since q is not a root of unity, we easily deduce from Equation (5.1) that $(g-h)/x^r \in \mathbb{C} \subset K$. Hence $g \in K$, as wanted.

In Case (b), we have that $a^* \notin q^{\mathbb{Z}}$ and $\tilde{b} = \rho(h) - a^*h$ for some $h \in K$. It follows that $\rho(g-h) = cx^\alpha(g-h)$, where $g-h \in F_0$ is a Puiseux series. Arguing as in Case (a), we obtain that $\alpha = 0$ and $g-h = c_0x^r$ for some $r \in \mathbb{Q}$ and $c_0 \in \mathbb{C}$. Hence $g \in K$. This ends the proof. \square

For order one equations over K whose solutions do not necessarily belong to F_0 , one has the following criterion for differential algebraicity.

Lemma 5.2. *Let L be a (ρ, δ) -field extension of \tilde{K} with $L^\rho = \tilde{C}$. Let $f \in L^*$ such that $\rho(f) = af$ with $a \in K^*$. Then, f is δ -algebraic over \tilde{K} if and only if $a = cx^\alpha \frac{\rho(g)}{g}$ for some $c \in \mathbb{C}^*$, $\alpha \in \mathbb{Q}$, and $g \in K^*$. Moreover, $\alpha = 0$ in Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{M} .*

Proof. Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{Q} can be deduced in a similar way to [HS08, Corollary 3.4] (resp. Case \mathbf{M} is given by [DHR18, Proposition 3.1]), combined with some descent argument to go from \tilde{C} to \mathbb{C} . Such argument is similar to the one given in the proof of [HS08, Corollary 3.2]. \square

5.2. Connected and irreducible Galois groups. As a second step, we consider the situation where the difference Galois group is both connected and irreducible.

Proposition 5.3. *Let us assume that $n \geq 2$. If $f \in F_0$ is a nonzero solution to Equation (1.2) and if the corresponding difference Galois group G over K is both connected and irreducible, then f is ∂ -transcendental over K .*

Proposition 5.3 is obtained as a direct consequence of the following more general result.

Proposition 5.4. *Let us consider a linear system of the form (4.1) with $n \geq 2$. Let us assume that the difference Galois group G for (4.1) over K is connected and irreducible. Let $\tilde{\mathcal{Q}}$ be a (ρ, δ) -Picard-Vessiot field extension for (4.1) over \tilde{K} , with fundamental matrix of solutions U . Then every column of U contains at least one element that is δ -transcendental over \tilde{K} .*

Proof of Proposition 5.3. We argue by contradiction, assuming that f is ∂ -algebraic over K . Then all coordinates of the vector $(f, \dots, \rho^{n-1}(f))^\top$ are also ∂ -algebraic over K . This follows from the fact that ρ and ∂ almost commute, that is $\partial\rho = c\rho\partial$ with $c = 1$ in Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{Q} , and $c = p$ in Case \mathbf{M} . We deduce from the construction of δ with respect to ∂ given in Section 3 (and the fact that \log is δ -algebraic for Case \mathbf{M}) that all coordinates of the vector $(f, \dots, \rho^{n-1}(f))^\top$ are also δ -algebraic over \tilde{K} .

Now, let A denote the companion matrix associated with Equation (1.2). Since $(f, \dots, \rho^{n-1}(f))^\top \in F_0^n$ is nonzero, Corollary 4.13 ensures the existence of a positive integer r and a (ρ^r, δ) -Picard-Vessiot field extension $\tilde{\mathcal{Q}}$ for $\rho^r(Y) = A_{[r]}Y$ over \tilde{K} such that the vector $(f, \dots, \rho^{n-1}(f))^\top$ is the first column of a fundamental matrix U . Furthermore, Remark 4.11 ensures that the Galois group of $\rho^r(Y) = A_{[r]}(Y)$ over K is equal to G , for the latter is connected. Thus, Proposition 5.4 applies with ρ replaced by ρ^r , providing a contradiction. \square

It remains to prove Proposition 5.4. As a key argument, we will use the following result due to Arreche and Singer [AS17, Lemma 5.1]. It says that the parametrized Galois group must be as big as possible when the difference Galois group has an identity component that is semisimple.

Proposition 5.5. *Let us consider a linear system $\rho(Y) = AY$, where $A \in \mathrm{GL}_n(\mathbb{C}(x))$ in Cases \mathbf{S}_0 and \mathbf{S}_∞ , and where $A \in \mathrm{GL}_n(\mathbb{C}(x^{1/\ell}))$ for some positive integer ℓ in Cases \mathbf{Q} and \mathbf{M} . In Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{M} , we let $G \subset \mathrm{GL}_n(\tilde{\mathcal{C}})$ denote the difference Galois group over \tilde{K} , and $H \subset \mathrm{GL}_n(\tilde{\mathcal{C}})$ denote the (ρ, δ) -Galois group over \tilde{K} . In Case \mathbf{Q} , we let $G \subset \mathrm{GL}_n(\tilde{\mathcal{C}})$ denote the difference Galois group over $\tilde{\mathcal{C}}(x^{1/\ell})$, and $H \subset \mathrm{GL}_n(\tilde{\mathcal{C}})$ denote the (ρ, δ) -Galois group over $\tilde{\mathcal{C}}(x^{1/\ell})$. As in Proposition 2.5, we see H as a subgroup of G . If the identity component of G is semisimple, then $H = G$.*

Proof. Cases \mathbf{S}_0 and \mathbf{S}_∞ are explicitly proved in [AS17, Lemma 5.1], since, by Remark 3.1, the field of differential constants of $\tilde{\mathcal{C}}$ is \mathbb{C} . Cases \mathbf{Q} and \mathbf{M} are not explicitly proved in [AS17, Lemma 5.1] for we consider the slightly more general situation where $A \in \mathrm{GL}_n(\mathbb{C}(x^{1/\ell}))$. However, the result follows easily from the argument given in the proof of [AS17, Lemma 5.1]. \square

We are now ready to prove Proposition 5.4.

Proof of Proposition 5.4. We let H denote the (ρ, δ) -Galois group of (4.1) over \tilde{K} , that is

$$H = \text{Gal}^\delta(\tilde{\mathcal{Q}}/\tilde{K}) \subset \text{GL}_n(\tilde{C}).$$

We also let \tilde{G} denote the Galois group of $\rho(Y) = AY$ over \tilde{K} . Let us argue by contradiction, assuming the existence of one column of U whose coordinates are all δ -algebraic over \tilde{K} .

We first show that all entries of U are δ -algebraic over \tilde{K} . Since G is irreducible, Lemma 4.6 implies that \tilde{G} is irreducible too. By Proposition 2.5, H is Zariski dense in the irreducible group \tilde{G} . This implies that H is irreducible, since otherwise, H would be conjugated to a group formed by block upper triangular matrices, as well as its Zariski closure, contradicting the irreducibility of \tilde{G} . Now, let \mathcal{S} denote the \tilde{C} -vector space of solution vectors over $\tilde{\mathcal{Q}}$ whose entries are all δ -algebraic over \tilde{K} . By assumption, \mathcal{S} is not reduced to $\{0\}$. Furthermore, \mathcal{S} is invariant under the action of any $\sigma \in H$. Since H is irreducible, we deduce that \mathcal{S} contains n linearly independent solutions to (4.1). In other words, all columns of U belong to \mathcal{S} , which implies that all entries of U are δ -algebraic over \tilde{K} , as claimed.

We observe that the determinant $\det(U)$ is solution to the equation

$$(5.2) \quad \rho(y) = \det(A)y$$

and that the difference Galois group of this equation over K is the group $\det(G)$. Since all entries of U are δ -algebraic over \tilde{K} , we obtain that $\det(U)$ is also δ -algebraic over \tilde{K} . We first consider the particular case where $\det(U) \in K$, and then we move to the general case.

Let us first assume that $\det(U)$ belongs to K . In that case, $\det(G) = \{1\}$ and therefore $G \subset \text{SL}_n(\mathbb{C})$. We recall that G is assumed to be connected and irreducible. According to Lemma 4.2, we have that G is primitive. By [SU93, Proposition 2.3], we finally obtain that G is semisimple. By Lemma 4.5, $\tilde{G} = G(\tilde{C})$ is semisimple too.

In Cases \mathbf{S}_0 , \mathbf{S}_∞ , and \mathbf{M} , we infer from Proposition 5.5 that $H = \tilde{G}$. Since all entries of U are δ -algebraic over \tilde{K} , the δ -dimension of the (ρ, δ) -Picard-Vessiot extension is zero^{||}, and therefore the δ -dimension of H is zero by [HS08, Proposition 6.26]. By a result of Kolchin [Kol73, Chap. IV, Proposition 10], the dimension of an algebraic group \mathcal{G} over \tilde{C} is the same as the δ -dimension of \mathcal{G} viewed as a differential group over \tilde{C} . This proves that the dimension of H equals the dimension of $G(\tilde{C})$. Since \mathbb{C} is algebraically closed, the dimension of $G(\tilde{C})$ over \tilde{C} equals the dimension of G over \mathbb{C} . Thus, the above equality $H = \tilde{G} = G(\tilde{C})$ implies that the dimension of the algebraic group G over \mathbb{C} is zero. We recall that an algebraic group of dimension zero is just a finite group, and that a finite connected group has cardinality one. Thus, we deduce that $G = \{I_n\}$, where we let I_n denote the identity matrix of size n . Since by assumption $n \geq 2$, this provides a contradiction with the assumption that G is irreducible. This ends the proof in these cases.

^{||}We refer the reader to [HS08, P. 374] for a definition of the notion of δ -dimension of a (ρ, δ) -Picard-Vessiot ring and of a (ρ, δ) -Galois group.

In Case **Q**, the system $\rho(Y) = AY$ has coefficients in $\mathbb{C}(x^{1/\ell})$ and the entries of U are δ -algebraic over $\tilde{C}(x^{1/\ell})$, for some positive integer ℓ . Let $G_{\tilde{K},\ell}$ denote the difference Galois group of $\rho(Y) = AY$ over $\tilde{C}(x^{1/\ell})$. Since \tilde{K} is an algebraic extension of $\tilde{C}(x^{1/\ell})$, we deduce from [Roq18, Theorem 7] that $G_{\tilde{K}}$ and $G_{\tilde{K},\ell}$ have the same identity component, which is $G_{\tilde{K}}$ since the latter is connected. Hence the identity component of the identity of $G_{\tilde{K},\ell}$ is semisimple. By Proposition 5.5, it follows that $H_\ell = G_{\tilde{K},\ell}$, where H_ℓ denote the (ρ, δ) -Galois group over $\tilde{C}(x^{1/\ell})$. Furthermore, the entries of U are δ -algebraic over $\tilde{C}(x^{1/\ell})$ which implies that the δ -dimension of H_ℓ is zero. Since $H_\ell = G_{\tilde{K},\ell}$, we get as previously that the algebraic group $G_{\tilde{K},\ell}$ has dimension zero. Since $G_{\tilde{K}} \subset G_{\tilde{K},\ell}$, this proves that the dimension of $G_{\tilde{K}}$ is also zero, and we can argue as previously to get a contradiction.

Now, let us consider the general case. We remind that $\det(U)$ is δ -algebraic over \tilde{K} and $\rho(\det(U)) = \det(A)\det(U)$. By Lemma 5.2, there exist some rational number α and nonzero elements $c \in \mathbb{C}$ and $g \in K$ such that $\det(A) = cx^\alpha \rho(g)/g$. Furthermore, $\alpha = 0$ in Cases **S**₀, **S** _{∞} , and **M**.

Let us consider the rank one linear difference system

$$(5.3) \quad \rho(y) = c^{-1/n} x^{-\alpha/n} y$$

Since $\tilde{\mathcal{Q}}$ is a (ρ, δ) -field with $\tilde{\mathcal{Q}}^\rho = \tilde{C}$, there exists a (ρ, δ) -Picard-Vessiot extension $\tilde{\mathcal{Q}}_1$ for (5.3) over $\tilde{\mathcal{Q}}$. Let $(\lambda) \in \mathrm{GL}_1(\tilde{\mathcal{Q}}_1)$ be a fundamental matrix associated with this system. Then

$$\rho(\lambda) = c^{-1/n} x^{-\alpha/n} \lambda$$

and λ is invertible in $\tilde{\mathcal{Q}}_1$. Using the commutativity of δ and ρ , we obtain that

$$\delta\left(\frac{\delta(\lambda)}{\lambda}\right) \in \tilde{\mathcal{Q}}_1^\rho = \tilde{C},$$

which shows that λ is δ -algebraic over \tilde{C} . In particular, it is δ -algebraic over \tilde{K} . Thus, all entries of the matrix $\lambda U \in \mathrm{GL}_n(\tilde{\mathcal{Q}}_1)$ are δ -algebraic over \tilde{K} .

On the other hand, we set

$$B = c^{-1/n} x^{-\alpha/n} A \in \mathrm{GL}_n(K) \quad \text{and} \quad U_B := \lambda U.$$

Note that $\rho(U_B) = BU_B$. Let G_B be the difference Galois group of the system $\rho(Y) = BY$ over K . Our choice of B ensures that $\det(B) = \rho(g)/g$. Thus, the equation

$$\rho(y) = \det(B)y$$

has a solution in K . It follows that its difference Galois group $\det(G_B)$ is reduced to $\{1\}$. Hence, $G_B \subset \mathrm{SL}_n(\mathbb{C})$. For $k \geq 1$, we let $G_{B_{[k]}}$ denote the difference Galois group of $\rho^k(Y) = B_{[k]}Y$ over K . By Proposition 4.10, there exists $r \geq 1$ such that $G_{B_{[r]}}$ is connected. Furthermore, $\rho^r(g)/g = \det(B_{[r]})$ and the difference Galois group of $\rho^r(Y) = B_{[r]}Y$ over K remains included in $\mathrm{SL}_n(\mathbb{C})$. Let us notice that

$$A_{[r]} = \alpha_r B_{[r]}$$

for some $\alpha_r \in K^\times$.

We claim that $G_{B_{[r]}}$ is irreducible. Let us argue by contradiction, assuming that $G_{B_{[r]}}$ is reducible. By Lemma 4.4, there exists $T \in \mathrm{GL}_n(K)$ such that $\rho^r(T)B_{[r]}T^{-1}$ is a block upper triangular matrix, and we deduce that

$$\alpha_r \rho^r(T)B_{[r]}T^{-1} = \rho^r(T)A_{[r]}T^{-1}.$$

By Lemma 4.4, we obtain that G_r , the difference Galois group of the system $\rho^r(Y) = A_{[r]}Y$ over K , is conjugated to a group of block upper triangular matrices, which implies that G_r is reducible. By Remark 4.11, $G_r = G$, providing a contradiction with the assumption that G is irreducible. This proves that $G_{B_{[r]}}$ is irreducible.

Finally by Lemma 4.2, the irreducible connected group $G_{B_{[r]}}$ is primitive. Again, since $G_{B_{[r]}} \subset \mathrm{SL}_n(\mathbb{C})$, we infer from [SU93, Proposition 2.3] that $G_{B_{[r]}}$ is semisimple. Since $\tilde{\mathcal{Q}}_1$ is a pseudo δ -field that contains \tilde{K} and all coordinates of U_B , we have $\tilde{K}\langle U_B \rangle \subset \tilde{\mathcal{Q}}_1$ (see Section 2.2.2 for the notation). Furthermore, since $\tilde{\mathcal{Q}}_1^\rho = \tilde{C}$, the pseudo δ -field $\tilde{K}\langle U_B \rangle$ is a (ρ^r, δ) -Picard-Vessiot extension for $\rho^r(Y) = B_{[r]}Y$ over \tilde{K} .

Since all entries of U_B are δ -algebraic over \tilde{K} , we can apply Proposition 5.5 and, arguing as in the first part, we deduce that $G_{B_{[r]}} = \{I_n\}$. Since $n \geq 2$, we obtain a contradiction with the fact that $G_{B_{[r]}}$ is irreducible. \square

5.3. The general case. We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2. We argue by induction on n . More precisely, our induction assumption reads as follows.

- (H _{n}) For all positive integers k and all $f \in F_0$ that is solution to a ρ^k -linear equation of order at most n with coefficients in K , we have either f is ∂ -transcendental over K or $f \in K$.

Proposition 5.1 implies that (H₁) hold true. Let $n \geq 2$ and let us assume (H _{$n-1$}). Let $f \in F_0$ be solution to a ρ^k -linear equation of order n with coefficients in K . Without any loss of generality, we can assume that $f \neq 0$ and $k = 1$. Considering the companion matrix associated with this equation, Corollary 4.13 ensures the existence of a positive integer r such that the following properties hold.

- (a) The vector $(f, \dots, \rho^{n-1}(f))^\top$ is solution to the system $\rho^r(Y) = AY$ for some $A \in \mathrm{GL}_n(K)$.
- (b) There exists a (ρ^r, δ) -Picard-Vessiot field extension $\tilde{\mathcal{Q}}$ for $\rho^r(Y) = AY$ over \tilde{K} , such that the vector $(f, \dots, \rho^{n-1}(f))^\top$ is the first column of a fundamental matrix $U \in \mathrm{GL}_n(\tilde{\mathcal{Q}})$.
- (c) The difference Galois group G of the system $\rho^r(Y) = AY$ over K is connected.

If G is irreducible, Proposition 5.3 shows that f is ∂ -transcendental. Hence (H _{n}) holds. From now on, we assume that G is reducible. Furthermore, we assume that f is ∂ -algebraic over K . Thus, it remains to prove that $f \in K$. Without loss of generality, we can assume that $r = 1$.

By Lemma 4.4, there exists a gauge transformation $T = (t_{i,j}) \in \mathrm{GL}_n(K)$ such that

$$\rho(T)AT^{-1} = \begin{pmatrix} A_1 & A_{1,2} \\ 0 & A_2 \end{pmatrix},$$

where $A_i \in \mathrm{GL}_{n_i}(K)$, $n_1 + n_2 = n$, and $n_2 < n$. Furthermore, let us assume that n_1 is minimal with respect to this property.

Set

$$(5.4) \quad (g_i)_{i \leq n}^\top := T(\rho^{i-1}(f))_{i \leq n}^\top \in F_0^n.$$

The vector $(g_i)_{n_1+1 \leq i \leq n}^\top \in F_0^{n_2}$ is solution to the system $\rho(Y) = A_2Y$. Furthermore, since f is ∂ -algebraic over K , the g_i are also ∂ -algebraic over K . By (H_{n_2}) and Remark 2.1, we obtain that

$$(5.5) \quad (g_i)_{n_1+1 \leq i \leq n}^\top \in K^{n_2}.$$

Let G_1 denote the difference Galois group of the system $\rho(Y) = A_1Y$ over K . We claim that G_1 is connected and irreducible. Indeed, if G_1 were reducible, then by Lemma 4.4, there would exist a gauge transformation changing A_1 into a block upper triangular matrix, contradicting the minimality of n_1 . Furthermore, the Galois group of $\rho(Y) = A_1Y$ over K is a quotient of the connected group G and thereby a connected group.

The main step of the proof consists in showing the following result.

Claim. One has $n_1 = 1$.

Proof of the claim. By assumption, f is ∂ -algebraic over K . Arguing as in the proof of Proposition 5.3, we get that all coordinates of the vector $(f, \dots, \rho^{n-1}(f))$ are δ -algebraic over \tilde{K} . Then, (5.4) implies that all the g_i 's are also δ -algebraic over \tilde{K} . We have to distinguish two cases.

Let us first assume that all the g_i 's belong to \tilde{K} . By (5.4) and Lemma 4.7, we obtain that

$$\forall i \in \{1, \dots, n\}, \quad g_i \in F_0 \cap \tilde{K} = K$$

and hence $(f, \dots, \rho^{n-1}(f))^\top = T^{-1}(g_1, \dots, g_n)^\top \in K^n$. Thus the vector $(f, \dots, \rho^{n-1}(f))^\top$ is fixed by the difference Galois group G . By Lemma 4.4, there exists $P \in \mathrm{GL}_n(K)$ such that

$$\rho(P)AP^{-1} = \begin{pmatrix} b_1 & B_{1,2} \\ 0 & B_2 \end{pmatrix},$$

for some matrices $B_{1,2}$ and B_2 with coefficients in K and a nonzero $b_1 \in K$. By minimality of n_1 , we obtain that $n_1 = 1$, as wanted.

Now, let us assume that at least one the g_i 's does not belong to \tilde{K} . Note that, by assumption, the f_i 's all belong to \tilde{Q} and thus the g_i 's all belong to \tilde{Q} too. We let H denote the (ρ, δ) -Galois group of $\rho(Y) = AY$ over \tilde{K} . By the parametrized Galois correspondence [HS08, Theorem 6.20], we deduce the existence of some $\sigma \in H$, such that

$$(\sigma(g_i))_{i \leq n}^\top \neq (g_i)_{i \leq n}^\top,$$

while (5.5) implies that $\sigma(g_i) = g_i$ for every i , $n_1 + 1 \leq i \leq n$. Set

$$u_1 := (g_i)_{i \leq n_1}^\top, \quad u_2 := (g_i)_{n_1+1 \leq i \leq n}^\top \quad \text{and} \quad v_1 := (\sigma(g_i))_{i \leq n_1}^\top.$$

Hence $w := u_1 - v_1$ is a nonzero vector. Since the coordinates of u_1 are δ -algebraic over \tilde{K} and σ belongs to the (ρ, δ) -Galois group H , the coordinates of v_1 are also δ -algebraic over \tilde{K} . Hence the coordinates of w are δ -algebraic over \tilde{K} . Furthermore, u_1 and v_1 are both solution to the system

$$\rho(Y) = A_1 Y + A_{1,2} u_2.$$

It follows that

$$\rho(w) = A_1 w.$$

Since we have

$$\tilde{C} = \tilde{K}^\rho \subset \tilde{K}(w)^\rho \subset \tilde{Q}^\rho = \tilde{C},$$

Corollary 4.13 ensures the existence of a positive integer s and a (ρ^s, δ) -Picard-Vessiot field extension \tilde{Q}_1 for the system $\rho^s(Y) = (A_1)_{[s]} Y$ over \tilde{K} such that w is the first column of a fundamental matrix. Furthermore, the difference Galois group of $\rho^s(Y) = (A_1)_{[s]} Y$ is equal to G_1 for the latter is connected. The coordinates of w being δ -algebraic over \tilde{K} and G_1 being connected and irreducible, Proposition 5.4 implies that $n_1 = 1$. \square

Now, let us prove that $(g_1, \dots, g_n) \in K^n$. Set

$$u_2 := (g_i)_{2 \leq i \leq n}^\top.$$

By (5.5), we have $u_2 \in K^{n-1}$ and thus $A_{1,2} u_2 \in K$. Furthermore, $g_1 \in F_0$ is solution to the inhomogeneous order one equation

$$\rho(Y) = A_1 Y + A_{1,2} u_2.$$

Since g_1 is ∂ -algebraic over K , Proposition 5.1 implies that $g_1 \in K$, and hence $(g_1, \dots, g_n) \in K^n$. By (5.4), we obtain that

$$(f, \dots, \rho^{n-1}(f))^\top = T^{-1}(g_1, \dots, g_n)^\top.$$

Since the coefficients of T belong to K , it follows that $f \in K$, as wanted. \square

REFERENCES

- [ADR18] C. E. Arreche, T. Dreyfus, and J. Roques. On the differential transcendence of the elliptic hypergeometric functions. *arXiv preprint arXiv:1809.05416*, 2018.
- [AF17] B. Adamczewski and C. Faverjon. Méthode de Mahler: relations linéaires, transcendance et applications aux nombres automatiques. *Proc. Lond. Math. Soc.*, 115:55–90, 2017.
- [AF18a] B. Adamczewski and C. Faverjon. Mahler’s method in several variables I: The theory of regular singular systems. *arXiv preprint arXiv:1809.04823*, 2018.
- [AF18b] B. Adamczewski and C. Faverjon. Mahler’s method in several variables II: Applications to base change problems and finite automata. *arXiv preprint arXiv:1809.04826*, 2018.
- [And01] Y. André. Différentielles non commutatives et théorie de Galois différentielle ou aux différences. *Ann. Sci. École Norm. Sup.*, 34:685–739, 2001.
- [AS17] C. E. Arreche and M. F. Singer. Galois groups for integrable and projectively integrable linear difference equations. *J. Algebra*, 480:423–449, 2017.
- [Béz94] J.-P. Bézivin. Sur une classe d’équations fonctionnelles non linéaires. *Funkcial. Ekvac.*, 37:263–271, 1994.
- [BG93] J.-P. Bézivin and F. Gramain. Solutions entières d’un système d’équations aux différences. *Ann. Inst. Fourier*, 43:791–814, 1993.
- [BM06] M. Bousquet-Mélou. Rational and algebraic series in combinatorial enumeration. In *International Congress of Mathematicians. Vol. III*, pages 789–826. Eur. Math. Soc., Zürich, 2006.

- [Bou90] N. Bourbaki. *Algebra. II. Chapters 4–7*. Elements of Mathematics. Springer-Verlag, Berlin, 1990.
- [Cas72] P. J. Cassidy. Differential algebraic groups. *Amer. J. Math.*, 94:891–954, 1972.
- [Cas89] P. J. Cassidy. The classification of the semisimple differential algebraic groups and the linear semisimple differential algebraic Lie algebras. *J. Algebra*, 121:169–238, 1989.
- [CH99] Z. Chatzidakis and E. Hrushovski. Model theory of difference fields. *Trans. Amer. Math. Soc.*, 351:2997–3071, 1999.
- [CHS08] Z. Chatzidakis, C. Hardouin, and M. F. Singer. On the definitions of difference Galois groups. In *Model Theory with Applications to Algebra and Analysis. Vol. 1*, volume 349 of *London Math. Soc. Lecture Note Ser.*, pages 73–109. Cambridge Univ. Press, Cambridge, 2008.
- [Coh65] R. M. Cohn. *Difference Algebra*. Interscience Publishers John Wiley & Sons, New York-London-Sydney, 1965.
- [CS07] P. J. Cassidy and M. F. Singer. Galois theory of parameterized differential equations and linear differential algebraic groups. In *Differential Equations and Quantum Groups*, volume 9 of *IRMA Lect. Math. Theor. Phys.*, pages 113–155. Eur. Math. Soc., Zürich, 2007.
- [DHR16] T. Dreyfus, C. Hardouin, and J. Roques. Functional relations of solutions of q -difference equations. *arXiv preprint arXiv:1603.06771*, 2016.
- [DHR18] T. Dreyfus, C. Hardouin, and J. Roques. Hypertranscendence of solutions of Mahler equations. *J. Eur. Math. Soc. (JEMS)*, 20:2209–2238, 2018.
- [DHRS18] T. Dreyfus, C. Hardouin, J. Roques, and M. F. Singer. On the nature of the generating series of walks in the quarter plane. *Invent. Math.*, 213:139–203, 2018.
- [DV12] L. Di Vizio. Approche galoisienne de la transcendance différentielle. In *Transcendance et Irrationalité*, volume 2012 of *SMF Journ. Annu.*, pages 1–20. Soc. Math. France, Paris, 2012.
- [Har08] C. Hardouin. Hypertranscendance des systèmes aux différences diagonaux. *Compos. Math.*, 144:565–581, 2008.
- [Har16] C. Hardouin. Galoisian approach to differential transcendence. In *Galois Theories of Linear Difference Equations: an Introduction*, volume 211 of *Math. Surveys Monogr.*, pages 43–102. Amer. Math. Soc., Providence, RI, 2016.
- [Hen97] P. A. Hendriks. An algorithm for computing a standard form for second-order linear q -difference equations. *J. Pure Appl. Algebra*, 117:331–352, 1997.
- [Hen98] P. A. Hendriks. An algorithm determining the difference galois group of second order linear difference equations. *J. Symbolic Comput.*, 26:445–462, 1998.
- [Hil02] D. Hilbert. Mathematical problems. *Bull. Amer. Math. Soc.*, 8:437–479, 1902.
- [Höl87] O. Hölder. Über die Eigenschaft der Gammafunction keiner algebraischen Differentialgleichung zu genügen. *Math. Ann.*, 28:1–13, 1887.
- [HS99] P. A. Hendriks and M. F. Singer. Solving difference equations in finite terms. *J. Symbolic Comput.*, 27:239–259, 1999.
- [HS08] C. Hardouin and M. F. Singer. Differential galois theory of linear difference equations. *Math. Ann.*, 342:333–377, 2008.
- [Ish98] K. Ishizaki. Hypertranscendence of meromorphic solutions of a linear functional equations. *Aequationes Math.*, 56:271–283, 1998.
- [Kol73] E. R. Kolchin. *Differential Algebra and Algebraic Groups*. Academic Press, New York-London, 1973. Pure and Applied Mathematics, Vol. 54.
- [KZ01] M. Kontsevich and D. Zagier. Periods. In *Mathematics Unlimited—2001 and Beyond*, pages 771–808. Springer, Berlin, 2001.
- [Mah29] K. Mahler. Arithmetische Eigenschaften der Lösungen einer Klasse von Funktionalgleichungen. *Math. Ann.*, 101:342–366, 1929.
- [Mah30a] K. Mahler. Arithmetische Eigenschaften einer Klasse transzendental-transzendenter Funktionen. *Math. Z.*, 32:545–585, 1930.
- [Mah30b] K. Mahler. Über das Verschwinden von Potenzreihen mehrerer Veränderlichen in speziellen Punktfolgen. *Math. Ann.*, 103:573–587, 1930.

- [Mil17] J. S. Milne. *Algebraic Groups. The Theory of Group Schemes of Finite Type over a Field*, volume 170. Cambridge University Press, 2017.
- [Moo96] E. H. Moore. Concerning transcendently transcendental functions. *Math. Ann.*, 48:49–74, 1896.
- [Ngu12] P. Nguyen. *Équations de Mahler et hypertranscendance*. PhD thesis, Institut de Mathématiques de Jussieu, 2012.
- [Nis84] Ke. Nishioka. A note on differentially algebraic solutions of first order linear difference equations. *Aequationes Math.*, 27:32–48, 1984.
- [Nis90] Ku. Nishioka. New approach in Mahler’s method. *J. Reine Angew. Math.*, 407:202–219, 1990.
- [OW15] A. Ovchinnikov and M. Wibmer. σ -Galois theory of linear difference equations. *Int. Math. Res. Not. IMRN*, 12:3962–4018, 2015.
- [Phi15] P. Philippon. Groupes de Galois et nombres automatiques. *J. Lond. Math. Soc.*, 92:596–614, 2015.
- [Ram92] J.-P. Ramis. About the growth of entire functions solutions of linear algebraic q -difference equations. *Ann. Fac. Sci. Toulouse Math.*, 1:53–94, 1992.
- [Ran92] B. Randé. Équations fonctionnelles de Mahler et applications aux suites p -régulières. *Thèse de l’Université Bordeaux 1*, 1992.
- [Roq18] J. Roques. On the algebraic relations between Mahler functions. *Trans. Amer. Math. Soc.*, 370:321–355, 2018.
- [Rub89] L. A. Rubel. A survey of transcendently transcendental functions. *Amer. Math. Monthly*, 96(9):777–788, 1989.
- [Sal63] R. Salem. *Algebraic Numbers and Fourier Analysis*. D. C. Heath and Co., Boston, Mass., 1963.
- [SS19] R. Schäfke and M. Singer. Consistent systems of linear differential and difference equations. *J. Eur. Math. Soc. (JEMS)*, 21:2751–2792, 2019.
- [SU93] M. F. Singer and F. Ulmer. Galois groups of second and third order linear differential equations. *J. Symbolic Comput.*, 16:9–36, 1993.
- [vdPS97] M. van der Put and M. F. Singer. *Galois Theory of Difference Equations*, volume 1666 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1997.
- [Wib10] M. Wibmer. *Geometric difference Galois theory*. PhD thesis, Heidelberg University, 2010.
- [Wib12] M. Wibmer. Existence of ∂ -parameterized Picard-Vessiot extensions over fields with algebraically closed constants. *J. Algebra*, 361:163–171, 2012.

UNIV LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1, CNRS UMR 5208, INSTITUT CAMILLE JORDAN, 43 BLVD. DU 11 NOVEMBRE 1918, F-69622 VILLEURBANNE CEDEX, FRANCE

E-mail address: `boris.adamczewski@math.cnrs.fr`

INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, UMR 7501 UNIVERSITÉ DE STRASBOURG ET CNRS, 7 RUE RENÉ DESCARTES 67084 STRASBOURG, FRANCE

E-mail address: `dreyfus@math.unistra.fr`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE ; UMR 5219, UNIVERSITÉ DE TOULOUSE ; CNRS, UPS, F-31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `hardouin@math.univ-toulouse.fr`